

1 Special Issue: The Multifunctional Fish Gill

2  
3 **Endocrine control of gill ionocyte function in euryhaline fishes**

4  
5 Jason P. Breves<sup>1,\*</sup> and Ciaran A. Shaughnessy<sup>2</sup>

6  
7 <sup>1</sup> *Department of Biology, Skidmore College, 815 N. Broadway, Saratoga Springs, NY*  
8 *12866, USA*

9 <sup>2</sup> *Department of Integrative Biology, Oklahoma State University, 501 Life Sciences West,*  
10 *Stillwater, OK 74078, USA*

11  
12 **\*Author for correspondence:**

13 Jason P. Breves, Ph.D.

14 Department of Biology

15 Skidmore College

16 815 N. Broadway

17 Saratoga Springs, NY 12866 USA

18 Phone: +1 518 580-5079

19 Fax: +1 518 580-5071

20 Email: jbreves@skidmore.edu

21  
22 **ORCID:**

23 J. Breves: 0000-0003-1193-4389

24 C. Shaughnessy: 0000-0003-2146-9126

25  
26 **Abstract:**

27 The endocrine system is an essential regulator of the osmoregulatory organs that  
28 enable euryhaline fishes to maintain hydromineral balance in a broad range of  
29 environmental salinities. Because branchial ionocytes are the primary site for the active  
30 exchange of Na<sup>+</sup>, Cl<sup>-</sup>, and Ca<sup>2+</sup> with the external environment, their functional regulation  
31 is inextricably linked with adaptive responses to changes in salinity. Here, we review the  
32 molecular-level processes that connect osmoregulatory hormones with branchial ion  
33 transport. We focus on how factors such as prolactin, growth hormone, cortisol, and  
34 insulin-like growth-factors operate through their cognate receptors to direct the

expression of specific ion transporters/channels,  $\text{Na}^+/\text{K}^+$ -ATPases, tight-junction proteins, and aquaporins in ion-absorptive (freshwater-type) and ion-secretory (seawater-type) ionocytes. While these connections have historically been deduced in teleost models, more recently, increased attention has been given to understanding the nature of these connections in basal lineages. We conclude our review by proposing areas for future investigation that aim to fill gaps in the collective understanding of how hormonal signaling underlies ionocyte-based processes.

**Keywords:** cortisol; growth hormone; ion transporter; prolactin; receptor; salinity

## 1. Introduction

Fishes, the most numerous and diverse vertebrates, consist of three major classes: Agnatha (jawless fishes), Chondrichthyes (cartilaginous fishes), and Osteichthyes (bony fishes) (Moyle and Cech 2004). Teleosts (class Osteichthyes; subclass Actinopterygii; infraclass Neopterygii; division Teleostei) and lampreys (members of class Agnatha) typically maintain extracellular fluids between 270 and 400 mOsm/kg, with  $\text{Na}^+$  and  $\text{Cl}^-$  constituting the major dissolved ions (Hwang and Lin 2014; Ferreira-Martins et al. 2016). Therefore, when residing in dilute freshwater (FW) environments, they are at risk for both excessive hydration and salt loss across body surfaces. To counterbalance this situation, the gill actively absorbs ions ( $\text{Na}^+$ ,  $\text{Cl}^-$ , and  $\text{Ca}^{2+}$ ) from the external environment, while the kidney and urinary bladder produce large volumes of dilute urine (Marshall and Grosell 2006; Kaneko et al. 2008). Lampreys and teleosts residing in seawater (SW), on the other hand, must excrete ions gained by passive diffusion from the surrounding environment and replace water that is lost via osmosis. While multiple segments of the gastrointestinal tract work in concert to promote solute-linked water absorption (Barany et al. 2020; Takei 2021), the gill and kidney secrete monovalent ( $\text{Na}^+$ ,  $\text{Cl}^-$ ) and divalent ( $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ , and  $\text{SO}_4^{2-}$ ) ions into the external environment, respectively (Kaneko et al. 2008). Cartilaginous fishes are typically marine in their distribution and operate as osmoconformers by retaining urea and trimethylamine oxide while maintaining internal  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations below those of SW (Hwang and Lin 2014). Hagfishes (members of class Agnatha) are marine osmoconformers with limited capacities to regulate internal ion concentrations.

While most fishes inhabit a single aquatic environment characterized as either FW ( $\leq 0.5\%$ ) or SW (30-40%), a relatively small percentage of species (~5%) are

considered “euryhaline” and can withstand both conditions (Schultz and McCormick 2013). Euryhaline species possess the capacity to rapidly modulate ion- and water-transporting activities within the gill, gastrointestinal tract, kidney, and urinary bladder following changes in salinity (Takei et al. 2014). In turn, they offer valuable opportunities to resolve how cellular and molecular processes within osmoregulatory organs enable fish to transition between environmental salinities. As the branchial exchange of ions with the external environment is critical for maintaining osmoregulatory balance, decades of focused investigations have pursued how “ionocytes”, cells specialized for  $\text{Na}^+$ ,  $\text{Cl}^-$ , and  $\text{Ca}^{2+}$  transport, operate relative to environmental salinity (Evans et al. 2005; Dymowska et al. 2012).

## **2. Molecular aspects of ionocyte function**

### **2.1 Freshwater-type ionocytes in teleosts**

Historically, various models have been put forth to explain how the branchial ionocytes of FW-acclimated fishes actively absorb ions against strong electrochemical gradients (Hwang and Lin 2014). The contrasting models of FW-type ionocytes reflect, in part, the evolution of different strategies for  $\text{Na}^+$  and  $\text{Cl}^-$  uptake across the teleost lineage (Dymowska et al. 2012; Takei et al. 2014; Yan and Hwang, 2019). For euryhaline teleosts, the most comprehensive models of FW-type ionocytes are derived from rainbow trout (*Oncorhynchus mykiss*), Mozambique tilapia (*Oreochromis mossambicus*), and Japanese medaka (*Oryzias latipes*) (Dymowska et al. 2012; Hsu et al. 2014; Inokuchi et al. 2022). For basal fishes, recent progress has been made in the development of FW-type ionocyte models for sea lamprey (*Petromyzon marinus*) (Ferreira-Martins et al. 2021). Without question, insights into how ionocytes operate in stenohaline zebrafish (*Danio rerio*) have supported progress in the euryhaline species listed above (Guh et al. 2015).

In FW-type ionocyte models for salmonids, largely conceived from findings in rainbow trout, two distinct subtypes absorb environmental  $\text{Na}^+$ ,  $\text{Cl}^-$ , and  $\text{Ca}^{2+}$ . In one subtype, termed peanut lectin agglutinin positive ( $\text{PNA}^+$ ) cells,  $\text{Na}^+/\text{H}^+$  exchangers 2 and 3 (Nhe2 and -3; Slc9a2 and -3), epithelial  $\text{Ca}^{2+}$  channel (ECaC), and an Slc26-family anion exchanger are expressed in the apical membrane.  $\text{Na}^+/\text{K}^+$ -ATPase (Nka) mediates the basolateral movement of  $\text{Na}^+$ , while an uncharacterized pathway allows for the exit of  $\text{Cl}^-$  (Ivanis et al. 2008; Dymowska et al. 2012). The other ionocyte subtype, termed  $\text{PNA}^-$  cells, expresses an apical  $\text{Na}^+$  channel, purported to be acid-sensing ion channel 4

(Asic4), along with apical H<sup>+</sup>-ATPase. Na<sup>+</sup>/HCO<sub>3</sub><sup>-</sup> cotransporter 1 (Nbce1; Slc4a4) and Nka are also expressed in PNA<sup>-</sup> cells to mediate the basolateral exit of Na<sup>+</sup> (Parks et al. 2007; Dymowska et al. 2014).

Like in trout, there are multiple FW-type ionocytes operating within the branchial epithelium of euryhaline Mozambique tilapia. “Type II” ionocytes express a Na<sup>+</sup>/Cl<sup>-</sup> cotransporter (Ncc) in the apical membrane to transport Na<sup>+</sup> and Cl<sup>-</sup> into the cell interior (Hiroi et al. 2008). This Ncc is denoted Ncc2 (Slc12a10) and is not a member of the “conventional” Ncc1 (Slc12a3) clade (Motoshima et al. 2023). Nka and Clc family Cl<sup>-</sup> channel 2c (Clc2c) support the basolateral transport of Na<sup>+</sup> and Cl<sup>-</sup> from the ionocyte interior into the blood plasma, respectively (Pérez-Ruis et al. 2015; Wang et al. 2015; Breves et al. 2017b). While Ncc2-expressing ionocytes operate in euryhaline and stenohaline species spanning teleost clades (Wang et al. 2009; Hsu et al. 2014; Inokuchi et al. 2017; Lema et al. 2018), they are conspicuously absent in salmonids (Hiroi and McCormick 2012). In tilapia, a second type of Na<sup>+</sup>-absorptive ionocyte which expresses Nka, coined “Type III” ionocytes, is characterized by the apical localization of Nhe3 (Hiroi et al. 2008). The density of Type III ionocytes (along with *nhe3* expression) increases in the gill filaments of tilapia exposed to low-Na<sup>+</sup> conditions (Inokuchi et al. 2008, 2009).

## 2.2 Freshwater-type ionocytes in basal fishes

In lampreys, two FW-adaptive ionocytes have been proposed to support ion uptake (Bartels and Potter 2004; Reis-Santos et al. 2008; Ferreira-Martins et al. 2021). These two ionocytes differ most notably in their expression of Nka and H<sup>+</sup>-ATPase. A “larval FW ionocyte” highly expresses H<sup>+</sup>-ATPase but shows low expression of Nka, whereas a “FW ionocyte” (observed in larvae as well as post-metamorphic and adult stages) strongly expresses both H<sup>+</sup>-ATPase and Nka. Branchial H<sup>+</sup>-ATPase E subunit (*atp6v1e*) expression markedly decreases when lamprey acclimate to elevated salinities (Reis-Santos et al. 2008; Ferreira-Martins et al. 2016). The ionoregulatory role of H<sup>+</sup>-ATPase in FW gills typically involves its co-expression with a pathway for the electrochemically neutral uptake of environmental Na<sup>+</sup>. The absorption of environmental Na<sup>+</sup> by lampreys appears to involve the epithelium Na<sup>+</sup> channel (ENaC) (Ferreira-Martins et al. 2016), while Ncc supports both Na<sup>+</sup> and Cl<sup>-</sup> uptake (Barany et al. 2021b). Accordingly, both ENaC and Ncc are highly expressed in the gills of FW-acclimated lamprey and exhibit decreased expression during SW acclimation, although which particular cell-types express these transporters has not been fully elucidated. The co-

involvement of an apical carbonic anhydrase-powered  $\text{Cl}^-/\text{HCO}_3^-$  exchanger and a basolateral  $\text{Cl}^-$ -channel in  $\text{Cl}^-$  uptake has also been proposed, but the molecular identities of these transporters are unresolved (Bartels and Potter 2004; Ferreira-Martins et al. 2021).

### 2.3 Seawater-type ionocytes in teleosts

Within the branchial epithelium of marine/SW-acclimated teleosts, SW-type ionocytes actively secrete excess  $\text{Na}^+$  and  $\text{Cl}^-$  into the environment. SW-type ionocytes express Nka and  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  cotransporter 1 (Nkcc1; Slc12a2) in the basolateral membrane to energize and facilitate the  $\text{Na}^+$ - and  $\text{K}^+$ -coupled passage of  $\text{Cl}^-$  from blood plasma into the cell interior (Marshall and Grosell 2006; Kaneko et al. 2008). The catalytic  $\alpha$ -subunit of the Nka enzyme contains binding sites for ATP,  $\text{Na}^+$ , and  $\text{K}^+$  (Geering 2008). Two distinct isoforms of the  $\alpha$ -subunit ( $\alpha 1a$  and  $\alpha 1b$ ) were identified in salmonids, first by Richards et al. (2003). In salmonids and cichlids, these isoforms have functional capacities exclusive to either FW ( $\alpha 1a$ ) or SW ( $\alpha 1b$ ), with branchial expression “switching” from one to the other during salinity transitions (Bystriansky et al. 2006; Nilsen et al. 2007; McCormick et al. 2009; Tipsmark et al. 2011; Dalziel et al. 2014). Apically located cystic fibrosis transmembrane conductance regulator 1 (Cftr1) enables  $\text{Cl}^-$  to exit SW-type ionocytes and to enter the external environment (Marshall and Grosell 2006). With Nkcc1 and Cftr1 forming the pathway for transcellular  $\text{Cl}^-$  excretion, tight-junction complexes composed of claudins (Cldns) between ionocytes and adjacent accessory cells provide the paracellular route for  $\text{Na}^+$  to exit the gill (Marshall and Grosell 2006; Tipsmark et al. 2008b; Bui and Kelly 2014). Attendant increases in branchial Nka, Nkcc1, and Cftr1 expression coincide with SW-acclimation. For this reason, all three ion transporters are widely employed as markers of branchial ion-secretory capacity.

### 2.4 Seawater-type ionocytes in basal fishes

The pathways for branchial  $\text{Cl}^-$  secretion are far less resolved in basal fishes than in teleosts. Cftr orthologs are present in the genomes of sturgeon, bichir, and coelacanth (Shaughnessy and Breves 2021), yet none of these orthologs have been functionally characterized. A single Cftr ortholog was identified in sea lamprey; however, *cftr* expression is low in all larval, juvenile, and adult tissues aside from intestine (Ren et al. 2015). Moreover, compared with human Cftr, lamprey Cftr exhibits limited  $\text{Cl}^-$

conductance and reduced activation by cAMP (Cui et al. 2019). Given the limited Cl<sup>-</sup> conductance of lamprey Cftr and the lack of a *cftr* transcriptional response to SW exposure (Shaughnessy et al. unpublished), it is questionable whether Cftr mediates the secretion of Cl<sup>-</sup> by lamprey ionocytes known to express Nka and Nkcc1 (Shaughnessy and McCormick 2020). A recent analysis of the updated inshore hagfish (*Eptatretus burgeri*) genome assembly (Yu et al. 2023; Marlétaz et al. 2023) indicates that a *cftr* ortholog may be absent in hagfishes altogether (Yamaguchi et al. 2023).

### 3. Hormones and ionocytes

The endocrine system has long been appreciated as a central player in the homeostatic regulation of salt and water balance in fishes. Perturbations in internal osmotic and ionic conditions caused by changes in environmental salinity elicit the secretion of hormones that modulate ion- and water-transport by key osmoregulatory organs. Because these regulatory connections are indispensable to maintaining hydromineral balance, there is no shortage of literature that discusses how hormones impact the osmoregulatory physiology of fishes at the organismal, organ, and cellular levels (Hirano 1986; McCormick 2001; Manzon 2002; Evans et al. 2005; Sakamoto and McCormick 2006; Takei and McCormick 2013; Takei et al. 2014). Therefore, in this review, we do not address all established hormonal actions within the gills of fishes; rather, we focus on how hormones control the molecular components of ionocytes. We focus on the regulatory connections identified in euryhaline species but, in several instances, reference stenohaline zebrafish for added context. An expansive collection of endocrine factors undeniably contributes to regulating branchial ionocytes (Evans et al. 2005; Takei et al. 2014); however, the identification of molecular endocrine targets is largely based on studies that focused upon the “classical” FW- and SW-adapting hormones in fishes, namely prolactin (Prl), growth hormone (Gh), and cortisol. While this review is heavily weighted toward describing the actions of these three hormones, we also highlight promising areas for future investigations into how additional endocrine factors regulate ionocytes.

### 4. Freshwater-adaptive endocrine control

#### 4.1 Prolactin

Euryhaline models, and most famously, mummichog (*Fundulus heteroclitus*), supported the discovery that pituitary hormones are key regulators of osmoregulatory

organs (Pickford and Atz 1957). Pickford (1953) and Burden (1956) reported that hypophysectomized mummichogs could not survive in FW, and that pituitary brei injections rescued them from death. Prl was subsequently identified as the pituitary factor that enables individuals to reside in dilute environments (Pickford and Phillips 1959). Over the succeeding decades, it was firmly established that through its highly conserved actions on teleost osmoregulatory organs, Prl stimulates a spectrum of activities befitting FW-acclimation (Loretz and Bern 1982; Hirano 1986; Manzon 2002; Sakamoto and McCormick 2006; Breves et al. 2014a, 2020). Accordingly, pituitary *prl* expression and plasma Prl levels rise when fish acclimate to low-salinity conditions (Lee et al. 2006; Hoshijima and Hirose 2007; Fuentes et al. 2010; Seale et al. 2012). The notion that ionocytes are targets of Prl signaling was supported decades ago by the observation that Prl influences ionocyte populations in Mozambique and Nile (*O. niloticus*) tilapia (Herndon et al. 1991; Pisam et al. 1993; Flik et al. 1994). With respect to directing ionoregulatory function, Zhou et al. (2003) showed that exogenous Prl stimulated ion uptake in rainbow trout branchial epithelium. Patterns of Prl binding and *prl receptor (prlr)* gene expression reported in both euryhaline and stenohaline FW species further associated Prl signaling with ionocytes (Dauder et al. 1990; Prunet and Auperin 1994; Weng et al. 1997; Rouzic et al. 2001; Santos et al. 2001; Lee et al. 2006; Huang et al. 2007; Fiol et al. 2009; Breves et al. 2013). Additionally, the Prlr was localized to branchial ionocytes of tilapia and sea bream (*Sparus aurata*) (Weng et al. 1997; Santos et al. 2001).

Only recently have investigations into the actions of Prl become unencumbered by a paucity of molecular tools to study FW-type ionocytes. For example, the characterization of tilapia Type II ionocytes by Hiroi et al. (2008) provided an opportunity to link Prl with a specific molecular pathway for ion uptake, particularly Ncc2. Prl enables hypophysectomized tilapia to recruit Ncc2-expressing ionocytes during FW acclimation, an activity that does not require systemic intermediaries (Breves et al. 2010c; Inokuchi et al. 2015; Watanabe et al. 2016) (Fig. 1). Prl similarly regulates branchial *ncc2* expression in euryhaline mummichog (Breves et al. 2022) and Japanese medaka (Bossus et al. 2017), as well as in stenohaline zebrafish (Breves et al. 2013). Activated Prl receptors (Prlrs) can modulate the transcription of target genes through JAK/STAT and ERK/MAPK signaling (Huang et al. 2007; Fiol et al. 2009; Chen et al. 2011). In medaka, Prl stimulates *ncc2* via STAT5 activation rather than through ERK- or AKT-dependent pathways (Bollinger et al. 2018). Since Clc2c is expressed within Ncc2-

expressing ionocytes to facilitate basolateral  $\text{Cl}^-$  movement (Pérez- Ruis et al. 2015; Wang et al. 2015), it is fitting that Prl coordinately promotes *clc2c* and *ncc2* expression (Breves et al. 2017b; Breves 2019) (Fig. 1). In contrast, branchial *clc3* expression in tilapia is not controlled by Prl (Tang and Lee 2011; Breves et al. 2017b).

The potential for Ncc-dependent pathways to operate in the osmoregulatory organs of cartilaginous and jawless fishes has recently received increased attention. In Japanese-banded houndshark (*Triakis scyllium*), a “conventional” *ncc1* (*slc12a3*) is expressed within a subpopulation of gill ionocytes, termed type-B cells, where its expression increases upon transfer from full-strength SW to 30% SW (Takabe et al. 2016). Given that elasmobranch genomes are devoid of Ncc2-encoding genes (Motoshima et al. 2023), Ncc1 may assume a role in branchial  $\text{Na}^+$  and  $\text{Cl}^-$  absorption in elasmobranchs. Similarly, the branchial expression of *ncca* (*ncc1*) in sea lamprey is attenuated during SW acclimation (Ferreira-Martins et al. 2016; Barany et al. 2021b). Given the Prlr expression in lamprey gills, the next step is to assess whether the recently found Prl participates in modulating *ncca* when lamprey transition between FW and marine environments (Gong et al. 2020).

In two lampreys (*P. marinus* and *Lethenteron reissneri*), the expression of gene transcripts encoding ENaC subunits increases under low- $\text{Na}^+$  conditions (Ferreira-Martins et al. 2016; Tseng et al. 2022). Thus, ENaC may provide a means for lampreys to absorb  $\text{Na}^+$  from FW; this strategy for  $\text{Na}^+$  absorption is absent in cartilaginous and ray-finned fishes (Ferreira-Martins et al. 2021). Curiously, branchial gene expression of an ENaC subunit, *scnn1a*, decreases when inshore hagfish experience high-salinity conditions (Yamaguchi et al. 2023). Despite hagfishes exhibiting ionoconformity, this response suggests that  $\text{Na}^+$  movement in the gill may be more complex than previously thought. To our knowledge, endocrine control of ENaC subunit expression has not been addressed in any cyclostome and, in an analogous fashion as *ncca*, should be probed for links to the Prlrs identified in hagfish and lamprey (Gong et al. 2020).

While branchial ionocytes leveraging Ncc operate in species across the three major fish lineages, they are not found within salmonids (Hiroi and McCormick 2012). In turn, an apically located  $\text{Cl}^-/\text{HCO}_3^-$  exchanger (Slc26a6) may provide a pathway for  $\text{Cl}^-$  absorption by PNA<sup>+</sup> ionocytes in rainbow trout and other salmonids (Boyle et al. 2015; Leguen et al. 2015). Branchial *slc26a6a2* is elevated in FW- versus SW-acclimated Atlantic salmon (Takvam et al. 2021) and is a transcriptional target of Prl signaling (Breves et al. unpublished). Therefore, Slc26a6a2 may constitute a pathway for Prl-



stimulated  $\text{Cl}^-$  uptake in species lacking Ncc-expressing ionocytes (Zhou et al. 2003). Because Leguen et al. (2015) reported *clc2* expression in trout ionocytes (putative PNA<sup>+</sup> cells), Prl-based control of salmonid *clc2* isoforms also warrants investigation. Studies of this nature will enable comparisons of Prl-Clc2 connectivity between species that do, and do not, leverage Ncc2-expressing ionocytes.

Within the PNA<sup>-</sup> ionocytes of trout, Nbce1 supports the absorption of environmental  $\text{Na}^+$  by cotransporting  $\text{Na}^+$  and  $\text{HCO}_3^-$  across the basolateral membrane (Parks et al. 2007; Leguen et al. 2015). The apical entry of  $\text{Na}^+$  into PNA<sup>-</sup> cells was proposed to occur via Asic4 through its electrochemical linkage to  $\text{H}^+$ -ATPase (Dymowska et al. 2014). Under this scenario, intracellular  $\text{HCO}_3^-$  is supplied by carbonic anhydrase (Parks et al. 2007). In tilapia, Nbce1 operates in the basolateral membrane of Ncc2-expressing ionocytes (Furukawa et al. 2011). To our knowledge, Nbce1, Asic4,  $\text{H}^+$ -ATPase, and carbonic anhydrase have not been associated with Prl signaling in trout or tilapia.

In addition to Type II ionocytes, a second type of  $\text{Na}^+$ -absorptive ionocyte in tilapia (Type III ionocytes) is characterized by the apical expression of Nhe3 (Hiroi et al. 2008). Prl promotes *nhe3* gene expression in tilapia gill filaments (Inokuchi et al. 2015; Watanabe et al. 2016) whereas it has no such effect in mummichog or zebrafish (Breves et al. 2013, 2022) (Fig. 1). Because salmonids express Nhe2 and -3 within PNA<sup>+</sup> ionocytes, they will prove key in resolving the extent to which Prl regulates Nhes among teleosts (Ivanis et al. 2008; Hiroi and McCormick 2012). Unfortunately, the lack of information on Nhes in lamprey ionocytes precludes consideration of a Prl-Nhe connection (Ferreira-Martins et al. 2021). Recent pharmacological experiments performed in zebrafish implicated  $\text{K}^+$ -dependent  $\text{Na}^+/\text{Ca}^{2+}$  exchangers (Nckxs) in mediating  $\text{Na}^+$  absorption (Clifford et al. 2022). Should roles emerge for Nckxs in supporting  $\text{Na}^+$  uptake by euryhaline species, Nckx isoforms would be additional candidates for regulation by Prl.

Nka plays a critical role in energizing ion transport by FW- and SW-type ionocytes, with the reciprocal expression of *nka- $\alpha$ 1a* and *- $\alpha$ 1b* first described in salmonids transitioning between FW and SW environments (Richards et al. 2003; Mackie et al. 2005; Bystriansky et al. 2006; Madsen et al. 2009; McCormick et al. 2009; Dalziel et al. 2014). Tilapia also undergo *nka- $\alpha$ 1a* and *- $\alpha$ 1b* “switching” upon salinity changes, and Prl stimulates the “FW-inducible” *nka- $\alpha$ 1a* isoform (Tipsmark et al. 2011; Breves et al. 2014b; Inokuchi et al. 2015; Watanabe et al. 2016) (Fig. 1). Thus far, the

capacity for Prl to promote *nka- $\alpha$ 1a* expression seems specific to tilapia, as Prl fails to stimulate *nka- $\alpha$ 1a* in Atlantic salmon (Tipsmark and Madsen 2009; Breves et al. unpublished). In zebrafish, *nka- $\alpha$ 1a1a.2* is expressed in Ncc2-expressing ionocytes responsible for Cl<sup>-</sup> uptake (Liao et al. 2009); however, Prl has no effect on branchial *nka- $\alpha$ 1a1a.2* expression (Breves 2019). The auxiliary  $\gamma$ -subunit of Nka (also called Fxyd) participates in the regulation of enzymatic activity by associating with the Na<sup>+</sup>/K<sup>+</sup> pump complex (Geering 2008; Pavlovic et al. 2013). Among the Fxyd isoforms identified in teleosts, Fxyd11 is predominately expressed in the gills where it interacts with Nka (Tipsmark 2008; Wang et al. 2008; Saito et al. 2010). In tilapia, Prl and cortisol synergistically promote *fxyd11* expression in FW (Tipsmark et al. 2011).

For teleosts residing in FW, greater than 90% of whole-body Ca<sup>2+</sup> uptake is mediated by branchial/epidermal ionocytes (Flik et al. 1996; Lin and Hwang 2016). Transcellular Ca<sup>2+</sup> uptake entails the apical entry of Ca<sup>2+</sup> through ECaC (Trpv5/6) followed by basolateral exit via Ca<sup>2+</sup>-ATPase 2 (Pmca2) and Na<sup>+</sup>/Ca<sup>2+</sup> exchanger 1 (Ncx1) (Flik et al. 1996; Liao et al. 2007). Prl is hypercalcemic in multiple teleosts (Pang et al. 1978; Fargher and McKeown 1989; Flik et al. 1989, 1994; Kaneko and Hirano 1993; Chakraborti and Mukherjee 1995; Wongdee and Charoenphandhu 2013), at least in part by stimulating branchial Pmca activity (Flik et al. 1996). Future investigations employing both euryhaline and stenohaline FW models are needed to determine whether Prl promotes ECaC and Ncx1 expression in parallel with promoting Pmca activity to sustain Ca<sup>2+</sup> uptake.

Aquaporins (Aqps) constitute a superfamily of integral membrane proteins that facilitate passive movements of water and small non-ionic compounds across cell membranes (Cerdà and Finn 2010). Multiple branchial cell types, including ionocytes, express a subset of Aqps (Lignot et al. 2002; Hirata et al. 2003; Watanabe et al. 2005; Tse et al. 2006; Brunelli et al. 2010; Tingaud-Sequeira et al. 2010; Tipsmark et al. 2010; Jung et al. 2012; Breves et al. 2016; Ruhr et al. 2020). Prl stimulates the expression of the aquaglyceroporin, Aqp3, in Mozambique tilapia (Breves et al. 2016) (Fig. 1), Japanese medaka (Ellis et al. 2019), and mummichog (Breves et al. 2022). On the other hand, Prl does not promote branchial *aqp1* expression (Ellis et al. 2019). Although the Aqp-specific effects of Prl suggest that Aqp3 plays an important role in FW-acclimated fish, there is still no clear picture of how it underlies adaptive processes. A role for Aqp3 in enhancing transepithelial water movement appears unlikely because branchial water exchange is disadvantageous to systemic hydromineral balance. Alternatively, Aqp3

may render ionocytes osmosensitive to extracellular conditions and/or capable of efficiently regulating their volume (Cutler and Cramb 2002; Watanabe et al. 2005; Tipsmark et al. 2010).

Prl has long been recognized for its effects on membrane permeability which result in a general “tightening” to minimize diffusive ion loss (Potts and Evans 1966; Hirano 1986). Paracellular solute movements across epithelia are governed in large part by the barrier properties of tight-junction complexes composed of Cldn and occludin proteins (Chasiotis et al. 2012). In tilapia and medaka, FW acclimation entails the increased expression of branchial *cldn28a* and *-28b*, respectively (Tipsmark et al. 2008a; Bossus et al. 2015). In Atlantic salmon and medaka, Prl stimulates *cldn28a* and *-28b* gene expression (Tipsmark et al. 2009; Bossus et al. 2017). Prl-Cldn28 connectivity thus provides a means to regulate tight-junction properties for minimizing ion loss in FW. *Occludin* expression is also correlated with environmental salinity (Chasiotis et al. 2009; Kumai et al. 2011; Whitehead et al. 2011), making it a good candidate for regulation by Prl; however, to our knowledge, this link has yet to be examined.

Teleosts express two separate Prlrs, denoted Prlr1 (Prlra) and -2 (Prlrb), that differ in their responses to salinity changes (Huang et al. 2007; Pierce et al. 2007; Fiol et al. 2009; Tomy et al. 2009; Rhee et al. 2010; Breves et al. 2011; Chen et al. 2011; Flores and Shrimpton 2012; Breves et al. 2013). Branchial *prlr1* has emerged as a transcriptional target of Prl in tilapia, mummichog, and zebrafish (Inokuchi et al. 2015; Breves et al. 2013, 2022). In turn, Prl seemingly upregulates the expression of Prlr1 to enhance the sensitivity of ionocytes to circulating hormone during FW acclimation (Weng et al. 1997). Alternatively, *prlr2/b* is typically insensitive to Prl (Breves et al. 2013, 2022; Inokuchi et al. 2015), which is not surprising given that its expression is upregulated by the hyperosmotic extracellular conditions associated with SW acclimation (Fiol et al. 2009; Inokuchi et al. 2015; Seale et al. 2019).

In tandem with initiating active ion uptake, euryhaline species must attenuate branchial ion secretion when transitioning from SW to FW. While promoting the recruitment of FW-type ionocytes and the expression of their associated ion transporters, Prl simultaneously dampens cellular and molecular phenotypes appropriate for SW conditions. For instance, Herndon et al. (1991) observed that Prl reduced the size and number of SW-type ionocytes in tilapia. At the molecular level, Prl inhibits the transcription of *nkcc1* and *cftr1* within the SW-type ionocytes of medaka and mummichog (Tipsmark and Madsen 2009; Bossus et al. 2017; Breves et al. 2022) (Fig.

2). Prl also inhibits branchial Nka activity and *nka- $\alpha$ 1b* expression (Pickford et al. 1970; Sakamoto et al. 1997; Shrimpton and McCormick 1998; Kelly et al. 1999; Mancera et al. 2002; Tipsmark and Madsen 2009), which, like *nkcc1* and *cftr1*, are elevated in SW to support ion secretion. Recall that while Cftr1 is the conduit for Cl<sup>-</sup> to exit SW-type ionocytes, tight junction complexes between ionocytes and accessory cells provide the paracellular path for Na<sup>+</sup> to exit the gill. The cation-selective tight-junctions adjacent to ionocytes are composed of multiple Cldn10 isoforms (Tipsmark et al. 2008b; Bui and Kelly 2014). Among the four mummichog *cldn10* genes (*cldn10c*, *-10d*, *-10e*, and *-10f*) upregulated in response to SW (Marshall et al. 2018), *cldn10f* is the only transcript downregulated by Prl (Breves et al. 2022) (Fig. 2). Collectively, these *nkcc1*, *cftr1*, and *cldn10f* responses illustrate the various means by which Prl inhibits branchial salt secretion.

#### 4.2 Growth hormone and somatolactin

As discussed in Section 5.1, Gh is conventionally regarded as a “SW-adapting hormone” because it promotes the survival of euryhaline fishes (and especially salmonids) in hyperosmotic environments (Björnsson 1997; McCormick et al. 2002; Takei et al. 2014). To our knowledge, there is no direct evidence that Gh plays a role in regulating FW-type ionocytes. Nonetheless, Gh receptors (Ghrs) are expressed in the gills of euryhaline species regardless of whether they are acclimated to FW or SW (Pierce et al. 2007; Poppinga et al. 2007; Breves et al. 2011; Link et al. 2010); therefore, Ghrs are at least present to mediate any direct regulatory connections between circulating Gh and FW-type ionocytes. It is certainly plausible that Gh may indirectly regulate FW-type ionocytes through the synthesis of insulin-like growth-factors (Igfs) (Reinecke et al. 1997; Berishvili et al. 2006; Reindl and Sheridan 2012). In fact, black-chinned tilapia (*Sarotherodon melanotheron*) exhibit enhanced *ghr* and *igf1* expression in the gill during FW acclimation (Link et al. 2010). Similarly, zebrafish exhibit elevated pituitary *gh* and branchial *ghr* (*ghra* and *-b*), *igf1a*, and *-2a* expression when challenged with ion-poor conditions (Hoshijima and Hirose 2007; Breves et al. unpublished). However, whether the Gh/Igf system supports the molecular responses of tilapia and zebrafish ionocytes to FW/ion-poor conditions has yet to be determined.

Somatolactin (SI), a member of the Gh/Prl-family of pituitary hormones, is a putative regulator of various physiological processes in fishes, particularly Ca<sup>2+</sup> homeostasis (Kaneko and Hirano 1993). Rainbow trout transferred to Ca<sup>2+</sup>-rich FW

exhibit reduced *sl* gene expression in the pituitary, a response that is consistent with SI having hypercalcemic activity (Kakizawa et al. 1993). Given the substantial progress made toward understanding how ionocytes absorb environmental  $\text{Ca}^{2+}$  (Lin and Hwang 2016), a reassessment of whether SI is indeed hypercalcemic is warranted by probing targets such as *ECaC*, *Pmca2*, and *Ncx1*.

#### 4.3 Cortisol

Cortisol is typically deemed a “SW-adapting hormone” because it directly stimulates the activities and/or expression of transporters tied to branchial ion-secretion (Section 5.2). The recognition that cortisol also promotes ion uptake in some teleosts arrived after its SW-adaptive role was firmly established (McCormick 2001; Takei and McCormick 2013). Morphological responses to cortisol in the gills of rainbow trout and American eel (*Anguilla rostrata*) suggested that FW-type ionocytes are targets of cortisol signaling (Perry et al. 1992), a notion that would be later supported with the development of molecular tools to more precisely study FW-type ionocytes. In tilapia, medaka, and zebrafish, *Nhe3* and *Ncc2* are expressed in distinct ionocyte subtypes (Hiroi and McCormick 2012; Hsu et al. 2014; Guh et al. 2015). In zebrafish, cortisol stimulates  $\text{Na}^+$  uptake in a fashion dependent upon the presence of *Nhe3b*-expressing ionocytes and promotes the differentiation of *Ncc2*-expressing ionocytes from a progenitor population (Kumai et al. 2012; Cruz et al. 2013a). While cortisol similarly promotes *ncc2* expression in medaka (Bossus et al. 2017; Ellis et al. 2019), this is not the case in tilapia (Breves et al. 2014b; Watanabe et al. 2016).

The FW-adaptive role of cortisol in zebrafish appears to be mediated solely by the glucocorticoid receptor (*Gr*) rather than the mineralocorticoid receptor (*Mr*) (Cruz et al. 2013b). The zebrafish *Gr* is expressed by *Nka*-rich branchial and epidermal ionocytes, with knockdown of *gr*, but not *mr*, disrupting the development of FW-type ionocytes through the action of forkheadbox I3 transcription factors (*Foxi3a* and *-b*) (Cruz et al. 2013b). Exogenous cortisol increases *nhe3b*, *H<sup>+</sup>-ATPase  $\alpha$ -subunit* (*atp6v1a*), and *ecac* expression in zebrafish embryos. In medaka embryos, knockdown of *gr2*, but not *gr1* or *mr*, decreases the total number of epidermal ionocytes (Trayer et al. 2013). Conversely, in FW-acclimated tilapia, it was suggested that the *Mr*, rather than the *Gr*, controls cortisol-mediated development of *Nka*-rich branchial ionocytes (Wu et al. 2023). Accordingly, *mr* expression occurs in ionocyte precursors/epidermal stem cells (Wu et al., 2023).

In Atlantic salmon, cortisol upregulates gene transcription and protein abundance of the “FW-inducible” Nka- $\alpha$ 1a isoform (Kiilerich et al. 2007b; McCormick et al. 2008, 2012; Tipsmark and Madsen 2009). Cortisol also upregulates the “SW-inducible” Nka- $\alpha$ 1b isoform (Kiilerich et al. 2007b; Tipsmark and Madsen 2009; Breves et al. 2024), and thus, the capacity of cortisol to increase the expression of both Nka- $\alpha$ 1a and - $\alpha$ 1b is indicative of its dual role in promoting FW- and SW-adaptive processes. While cortisol was shown to stimulate branchial carbonic anhydrase activity in trout (Gilmour et al. 2011), to our knowledge, no ion transporters expressed in salmonid FW-type ionocytes outside of Nka (e.g., Nhe2, -3, Asic4, ECaC, and Nbce1) have been linked with cortisol. This is a significant knowledge gap, especially given that cortisol is known to stimulate  $\text{Ca}^{2+}$  uptake by ECaC-expressing ionocytes in zebrafish (Lin and Hwang 2016). Reminiscent of the scenario for Prl (Section 4.1), future work is warranted to resolve whether cortisol affects  $\text{Ca}^{2+}$  uptake pathways in euryhaline species.

In addition to promoting key mediators of ion uptake (e.g., Ncc2, Nhe3, and Nka- $\alpha$ 1a), cortisol promotes FW acclimation by decreasing the paracellular permeability of the branchial epithelium (Kelly and Wood 2002; Zhou et al. 2003; Kolosov and Kelly 2017). This important contribution to FW acclimation is achieved through the regulation of specific tight-junction proteins. For instance, cortisol increases the expression of *cldn8d*, *-10c*, *-10d*, *-10e*, *-10f*, *-11a*, *-27a*, *-30c*, and *-33b* in various euryhaline species (Tipsmark et al. 2009; Bui et al. 2010; Bossus et al. 2017; Kolosov and Kelly 2017). Finally, it certainly must be recognized that cortisol can promote FW acclimation by acting in concert with Prl (Eckert et al. 2001; McCormick 2001). For instance, from a molecular perspective, Prl and cortisol act synergistically to promote branchial *nka- $\alpha$ 1a* and *cldn28b* expression in tilapia and medaka, respectively (Watanabe et al. 2016; Bossus et al. 2017).

#### 4.4 Thyroid hormones

Although limited, there is evidence that thyroid hormones are involved in the control of FW-adaptive branchial processes. Unfortunately, information is particularly scant regarding plasma thyroxine ( $\text{T}_4$ ) and 3-3'-5-triiodothyronine ( $\text{T}_3$ ) levels in euryhaline species undergoing FW acclimation. In sea bream, plasma  $\text{T}_4$  levels increase following transfer from SW to FW (Klaren et al. 2007). Alternatively, Mozambique tilapia acclimating to FW exhibit rapid declines in both plasma  $\text{T}_4$  and  $\text{T}_3$  (Seale et al. 2021). While the dynamics of  $\text{T}_4$  and  $\text{T}_3$  in tilapia suggest a hyposmotically-induced suppression

of thyroid hormone production at the systemic level, at the level of the gill, these changes coincide with an increase in the outer-ring deiodination activity of deiodinase 2 (Dio2). As shown in mummichog, Dio2 expression/activity is activated by hyposmotic stress (López-Bojórquez et al. 2007). Thus, increased branchial Dio2 activity supports the local production of T<sub>3</sub> at a time when the recruitment of ionocytes is activated following entry into FW (Hiroi et al. 2008; Breves et al. 2021). Accordingly, tilapia treated with T<sub>4</sub> exhibit an increase in the density and size of presumed FW-type ionocytes (Peter et al. 2000). It remains to be seen whether these cellular responses to T<sub>4</sub> manifest changes in branchial *ncc2*, *nhe3*, and *clc2c* expression.

## **5. Seawater-adaptive endocrine control**

### **5.1 Growth hormone and insulin-like growth-factors**

Although much of the early attention given to the Gh/Igf system in fishes was driven by its potential application to understanding growth in aquaculture settings, the osmoregulatory actions of both Gh and Igf1 have emerged as important aspects of the hormonal control of osmoregulation. In salmonids, Gh is integral to the timing of parr-smolt transformation and the associated development of SW tolerance (Hoar 1988; Björnsson 1997; McCormick 2013), and accordingly, plasma Gh levels increase during smolting (Boeuf et al. 1989; Prunet et al. 1989; Young et al. 1989; McCormick et al. 2007, 2013; Nilsen et al. 2008). The SW-adaptive role for Gh is not restricted to salmonids, as in both salmonid and non-salmonid teleost species, exposure to SW corresponds with elevated plasma Gh levels alongside with increased *gh* gene expression, Gh protein content, and somatotroph numbers in the pituitary (Deane and Woo 2009). As shown in Mozambique tilapia, somatotrophs release Gh in direct response to hyperosmotic extracellular conditions (Seale et al. 2002). Importantly, treatment with Gh upregulates branchial Nka activity and improves the SW tolerance of several euryhaline teleosts (Madsen 1990a, b; McCormick 1996; Xu et al. 1997; Mancera and McCormick 1998; Pelis and McCormick 2001). Intraperitoneal injection with Gh also increases Nkcc1 protein abundance within SW-type ionocytes (Pelis and McCormick 2001) and stimulates *nka-α1b* and *nkcc1* expression (Tipsmark and Madsen 2009), although these effects were most pronounced when Gh was co-administered with cortisol.

Ghrs are expressed in teleost gills (Gray et al. 1990; Yao et al. 1991; Sakamoto and Hirano 1991); however, they have yet to be localized to any discrete branchial cell-

types. It was initially reported that rainbow trout acclimating to SW do not exhibit changes in branchial Gh binding (Sakamoto and Hirano 1991). More recent molecular analyses describe variable branchial *ghr* expression patterns with respect to SW acclimation. In Atlantic salmon, *ghr* expression has been seen to increase (Kiilerich et al. 2007a; Nilsen et al. 2008) or not change at all (Breves et al. 2017a) during smolting. Likewise, there is little consistency in branchial *ghr* patterns following SW exposure, with increases, decreases, and no changes in expression all having been observed across several species (Kiilerich et al. 2007a; Nilsen et al. 2008; Breves et al. 2010a, b; Flores and Shrimpton 2012; Einarsdóttir et al. 2014; Breves et al. 2017a; Link et al. 2022). Additionally, Gh-treated gill explants from coho salmon (*Oncorhynchus kisutch*) and Nile tilapia did not exhibit changes in Nka activity, or *nka-α1b* and *nkcc1* gene expression (McCormick et al. 1991; Breves et al. 2014b). Rather than directly regulating the expression of specific ion-transporters, Gh may exert cytogenic effects that promote the recruitment of branchial ionocytes (Madsen 1990a, b; Flik et al. 1993; Prunet et al. 1994). For instance, Gh-elicited increases in Nka activity and *Nkcc1* in Atlantic salmon were coincident with an increased abundance of ionocytes (Pelis and McCormick 2001).

Gh is the primary regulator of the production and release of Igf1 and -2 from the liver (Pierce et al. 2011; Reindl and Sheridan 2012). Branchial *igf1 receptor (igf1r)* expression increases during smolting and upon exposure to SW (Nilsen et al. 2008; Shimomura et al. 2012), and increased circulating Igf1 levels correlate with elevated branchial Nka activity (Agustsson et al. 2001; McCormick et al. 2007; Shimomura et al. 2012). However, not all studies have observed rises in plasma Igf1 during smolting (Nilsen et al. 2008; Breves et al. 2017a). Intraperitoneal injection of Atlantic salmon with Igf1 increases SW tolerance but only marginally impacts gill Nka activity (McCormick 1996) whereas *Nkcc1* in isolated Japanese eel (*Anguilla japonica*) gill cells is stimulated by Igf1 (Tse et al. 2007). In addition to exerting osmoregulatory actions as endocrine signals (i.e., secreted from the liver and acting upon ionocytes) (Madsen and Bern 1993), Igf1 and -2 may also operate as autocrine/paracrine signals (i.e., produced by and acting upon ionocytes) (Berishvili et al. 2006; Tipsmark and Madsen 2009). In Atlantic salmon, Nilsen et al. (2008) reported increases in gill *igf1* and *igf1r* during smolting and SW acclimation, even when no increase in circulating Igf1 was detected. Similarly, Breves et al. (2017a) observed increases in branchial *igf2* and *igf1ra* expression in smolts following SW exposure.



The promotion of SW-adaptive ionoregulatory capacities by Gh may be best explained by its interaction with cortisol to promote both the proliferation of ionocytes and their responsiveness to cortisol (McCormick 2013). Studies using salmonids demonstrated that cortisol interacts with the Gh/Igf system to affect SW-type ionocytes. The co-administration of cortisol with either Gh or Igf1 increases gill Nka activity to levels beyond those induced by treatment with each hormone individually (Madsen 1990a, b; Madsen and Korsgaard 1991; McCormick 1996). Scenarios proposed to underlie the apparent synergistic actions of cortisol and Gh include, 1) Gh promotes Gr abundance in ionocytes, thereby increasing the capacity for cortisol to affect ion transporter expression, and 2) Gh promotes ionocyte proliferation while cortisol promotes the differentiation of ionocytes (McCormick 2013). Thus, future work should leverage recent insights into the regulators of ionocyte differentiation, such as forkhead box transcription factors (Hsiao et al. 2007), to elucidate how Gh and cortisol shape SW-type ionocyte populations.

Recent studies also describe the potential for Gh and Igf1 to regulate SW-adaptive branchial processes in lampreys. Kawauchi et al. (2002) were the first to identify a lamprey Gh capable of stimulating hepatic *igf1* expression. Later, Gh-like cells in the lamprey pituitary were shown to increase in abundance during metamorphosis (Nozaki et al. 2008). Discovery of the Ghr, Prlr, and Prl itself in sea lamprey spurred recent investigations into their regulatory roles (Gong et al. 2022). Although pituitary *gh* and *prl* expression are upregulated during sea lamprey metamorphosis (Gong et al. 2022), it was later shown that *gh* also increases in the pituitary of non-metamorphosing larvae over the same period (Ferreira-Martins et al. 2023). Thus, such increases in *gh* expression may be seasonal, and it remains unclear whether the same is true for pituitary *prl* expression. In any case, branchial *ghr* and *prlr* gene expression also increases during metamorphosis (Gong et al. 2020; Ferreira-Martins et al. 2023). Because similar increases do not occur in non-metamorphosing larval lamprey (Ferreira-Martins et al. 2023), heightened *ghr* and *prlr* expression likely underlies developmental (as opposed to seasonal) processes. Substantial increases in hepatic and branchial *igf1* expression also occur throughout metamorphosis, and therefore, endocrine as well as autocrine/paracrine actions of Igf1 may operate in lamprey (Ferreira-Martins et al. 2023). Surprisingly, SW exposure does not affect pituitary *gh* expression, hepatic *igf* expression, or branchial *ghr* and *igf1* expression (Gong et al. 2020, 2022; Ferreira-Martins et al. 2023) and treatment with recombinant Gh does not affect branchial ion

transporters (Gong et al. 2022). Future studies in lamprey are warranted to assess whether Gh and Igf1 promote the recruitment of SW-type ionocytes through cytogenic actions.

## 5.2 Corticosteroids

In lobe-finned fishes (Sarcopterygii) and tetrapods, cortisol (or, in some cases, corticosterone) and aldosterone are the products of the corticosteroid biosynthesis pathway and the predominant circulating hormones. Cortisol and aldosterone separately regulate carbohydrate metabolism and osmoregulation by interacting with the Gr and Mr, respectively. In all other fishes, corticosteroids and their receptors mediate both carbohydrate metabolism and osmoregulation. However, important differences exist between fish groups, particularly with respect to the milieu of corticosteroids in circulation and the identity and expression of receptors that mediate their actions. Here, we focus on corticosteroids that are known to directly regulate branchial processes in fishes.

Non-sarcopterygian fishes lack aldosterone synthase (Cyp11b2) and consequently the ability to synthesize aldosterone (Baker 2003; Takahashi and Sakamoto 2013). In actinopterygian fishes, cortisol is the predominant corticosteroid present in circulation, with 11-deoxycorticosterone and corticosterone present at far lower concentrations (Prunet et al. 2006). Among the circulating corticosteroids in actinopterygians, cortisol has both glucocorticoid and mineralocorticoid activity. To a far lesser extent, 11-deoxycorticosterone also exhibits mineralocorticoid-like actions (Takahashi and Sakamoto 2013). Chondrichthyan fishes produce a novel steroid biosynthetic product, 1 $\alpha$ -hydroxycorticosterone, which exhibits some mineralocorticoid-like action (Anderson 2012). However, chondrichthyans do not utilize branchial processes for bulk ion secretion but rather use the salt-secretory rectal gland (Wright and Wood 2015); therefore, the potential ionoregulatory actions of 1 $\alpha$ -hydroxycorticosterone will not be discussed here. Lampreys apparently lack 11 $\beta$ -hydroxylase (Cyp11b1) and cannot produce cortisol or corticosterone (Bridgham et al. 2006; Close et al. 2010; Rai et al. 2015). Thus, 11-deoxycortisol and 11-deoxycorticosterone are the most abundant circulating corticosteroids in lampreys and exhibit capacities to regulate branchial ionoregulatory activities (Close et al. 2010; Shaughnessy et al. 2020).

Chondrichthyan and actinopterygian fishes express both classes of corticosteroid receptors (Gr and Mr). In actinopterygians, it has long been held that the ionoregulatory actions of corticosteroids result from cortisol acting through the Gr. While this remains true, recent discoveries have added some nuance to this perspective. For instance, particular teleosts express two distinct Gr orthologs (Bury et al. 2003) as well as an Mr (Colombe et al. 2000). Knowledge of these three corticosteroid receptor subtypes has motivated investigations into how the actions of cortisol and 11-deoxycorticosterone are differentially mediated by these receptors (see below). Interestingly, lamprey do not express Gr or Mr but rather an ancestral “corticoid receptor” (Cr) that facilitates the osmoregulatory actions of 11-deoxycortisol (Bridgham et al. 2006; Close et al. 2010; Shaughnessy et al. 2020).

Using adult sea lamprey, Close et al. (2010) demonstrated that 11-deoxycortisol elicits an increase in branchial Nka activity. Later, Shaughnessy et al. (2020) described how 11-deoxycortisol supports the acquisition of SW tolerance during metamorphosis. Plasma 11-deoxycortisol levels and gill Cr abundance both increase during metamorphosis and are positively correlated with gill Nka activity. Accordingly, the treatment of mid-metamorphic lamprey with 11-deoxycortisol improves SW tolerance and increases gill Nka and *Nkcc1* protein expression (Shaughnessy et al. 2020; Barany et al. 2021a). Likewise, 11-deoxycortisol increases the expression of *nka* and *nkcc1* transcripts in gill explants (Shaughnessy et al. 2020). Interestingly, 11-deoxycorticosterone can elicit modest increases in branchial *nka* and *nkcc1* expression but is far less potent than 11-deoxycortisol (Shaughnessy et al. 2020). Future studies are warranted to further elucidate the ionoregulatory roles of 11-deoxycortisol and 11-deoxycorticosterone, and particularly whether they interact with Gh and Prl.

Cortisol has long been known to support the acclimation of teleosts to SW. Multiple lines of evidence have described this role, including early studies demonstrating that circulating cortisol increases during salmonid parr-smolt transformation and upon exposure to SW (Fontaine and Hatey 1954; Specker and Schreck 1982; Langhorne and Simpson 1986; Shrimpton et al. 1994), and that SW tolerance is increased following cortisol treatment (Bisbal and Specker 1991). Elevations in plasma cortisol following exposure to SW also occur in numerous non-salmonid species (McCormick 2001). Early work described the direct action of cortisol to increase gill Nka activity, which correlated with the development of SW tolerance during smolting (Langhorne and Simpson 1986; McCormick and Saunders 1987). Additional studies showed that gill Nka activity can be

impacted *in vivo* by cortisol injections (Pickford et al. 1970; Bisbal and Specker 1991; McCormick et al. 1991) and *in vitro* by exposing gill explants to cortisol-containing media (McCormick and Bern 1989).

More recently, cortisol was shown to regulate proteins and gene transcripts expressed by SW-type ionocytes, such as Nka, Nkcc1, and Cftr (Fig. 3). Atlantic salmon interperitoneally injected with cortisol increase the expression of *nka- $\alpha$ 1b* (McCormick et al. 2008; Tipsmark and Madsen 2009; Breves et al. 2020, 2024) and the protein abundance of Nka and Nkcc1 (Pelis and McCormick 2001). In gill explants from FW- and SW-acclimated Atlantic salmon, cortisol increases *nka- $\alpha$ 1b* and *nkcc1* expression (Tipsmark et al. 2002; Kiilerich et al. 2007b, 2011a, b, c). *In vivo* treatment with cortisol increases *cftr1* expression in Atlantic salmon parr and smolts (Singer et al. 2003; Breves et al. 2020, 2024), and *in vitro* exposure of gill explants to cortisol increases *cftr1* and *nkcc1* (Kiilerich et al. 2007b). Likewise, cortisol promotes *cftr1* and *nkcc1* expression in the gills of FW-acclimated trout and medaka (Tipsmark et al. 2002; Kiilerich et al. 2011a; Bossus et al. 2017). In tilapia and striped bass (*Morone saxatilis*), cortisol similarly promotes branchial *nkcc1* expression (Kiilerich et al. 2011c). Cortisol also promotes components of SW-type ionocytes in non-teleost models, such as Nka and Nkcc1 in Atlantic and Persian sturgeon (*Acipenser oxyrhynchus* and *A. persicus*) (Khodabandeh et al. 2009; McCormick et al. 2020).

Fewer studies have examined the molecular actions of 11-deoxycorticosterone, as it circulates at far lower concentrations than cortisol. Intraperitoneal injection of 11-deoxycorticosterone has no effect on SW tolerance or branchial *nka- $\alpha$ 1a* and *- $\alpha$ 1b* expression in Atlantic salmon (McCormick et al. 2008). The *in vitro* effects of 11-deoxycorticosterone vary depending on whether treated filaments are collected from salmon acclimated to either FW or SW. 11-deoxycorticosterone is more effective in stimulating *nka- $\alpha$ 1a* versus *- $\alpha$ 1b* expression (Kiilerich et al. 2007b, 2011a, b), although this effect is generally far less consistent than that of cortisol.

The role of the Gr in mediating the ionoregulatory actions of cortisol in teleosts has also received considerable attention. Early studies demonstrated that a corticosteroid receptor expressed in the gills with high binding affinity for cortisol increases during parr-smolt transformation and SW acclimation (Weisbart et al. 1987; Maule and Schreck 1990; Shrimpton and Randall 1994; Shrimpton et al. 1994; Marsigliante et al. 2000). Moreover, Gr expression is strongly correlated with the capacity for cortisol to stimulate branchial Nka activity (Shrimpton and McCormick 1999).

Following the discovery of two distinct Grs (Bury et al. 2003) and an Mr (Colombe et al. 2000; Sturm et al. 2005) in teleost fishes, studies using selective receptor antagonists investigated their individual roles in mediating the actions of cortisol and 11-deoxycorticosterone. It was proposed that the Gr and Mr underlie the duality of cortisol operating as a FW- and SW-adapting hormone (Prunet et al. 2006). In support of this, the upregulation of *gr* expression occurs in the gills of several species during smolting or following SW exposure (Mazurais et al. 1998; Mizuno et al. 2001; Kiilerich et al. 2007a; Nilsen et al. 2008; Yada et al. 2014; Bernard et al. 2020), and a potential role for the Mr in FW ionoregulation has been suggested (Sloman et al. 2001; Scott et al. 2005; Kiilerich et al. 2011a). The ionoregulatory role of the Mr in FW may entail activation by both cortisol and 11-deoxycorticosterone, as the Mr is potently activated by both hormones (Sturm et al. 2005; Katsu et al. 2018). Investigations into the regulation of *gr* and *mr* during smolting or SW acclimation have generally presented mixed results. In some studies, only the *gr* is upregulated during smolting (Kiilerich et al. 2007a, 2011b; Nilsen et al. 2008), and in others, the transcriptional upregulation of both receptors occurred (Yada et al. 2014; Bernard et al. 2020). Similarly, there seems to be little consistency in how *gr* and *mr* are transcriptionally regulated during SW acclimation in salmonids (Kiilerich et al. 2007b, 2011a; Nilsen et al. 2008; Flores and Shrimpton 2012) as well as in non-salmonids (Aruna et al. 2012a, b).

Several *in vivo* and *in vitro* studies have employed receptor blockade approaches, including the cotreatment of corticosteroids with mammalian Gr and Mr antagonists (e.g., RU486 and spironolactone, respectively). Cotreatment with RU486 blocks the upregulation of branchial *nka- $\alpha$ 1a* and *- $\alpha$ 1b* by cortisol, whereas cotreatment with spironolactone has no effect on SW tolerance or *nka- $\alpha$ 1a* and *- $\alpha$ 1b* expression (McCormick et al. 2008). Kiilerich et al. (2007b) demonstrated using Atlantic salmon gill explants that both RU486 and spironolactone can block the ability of cortisol to upregulate *nka- $\alpha$ 1a*, *- $\alpha$ 1b*, and *cfr1*. However, these results were not consistent across species or salinities (Kiilerich et al. 2007b, 2011b, c). In teleosts, RU486 antagonizes both Gr1 and -2, with more potent effects on Gr1 (Bury et al. 2003). On the other hand, spironolactone is now known to agonize the fish Mr, activating it with similar potency as cortisol, 11-deoxycorticosterone, and aldosterone (Sugimoto et al. 2016; Fuller et al. 2019). Thus, studies which use RU486 and spironolactone to differentially block the Gr and Mr should be interpreted with caution. Considering the challenges associated with pharmacologically targeting the fish Gr and Mr, advanced molecular approaches using

transcriptional knockdown or transgenic knockout have emerged to investigate the Gr and Mr (Faught and Vijayan 2018; Yan and Hwang 2019). To date, these approaches have mostly been leveraged to investigate the metabolic, developmental, and ionoregulatory actions of corticosteroids in zebrafish (Faught and Vijayan 2018; Yan and Hwang 2019), which cannot tolerate SW. However, Japanese medaka offer a promising euryhaline model for knockdown or knockout approaches (Yan and Hwang 2019) and is therefore poised to delineate the Gr- and Mr-mediated actions of corticosteroids on SW-type ionocytes.

In tetrapods, the interaction of aldosterone with the Mr is facilitated by coexpression of the Mr with the cortisol-inactivating enzyme, 11 $\beta$ -hydroxylase 2 (Cyp11b2). Interestingly, a strong transcriptional upregulation of *cyp11b2* occurs in the gills of smolting Atlantic salmon (Kiillerich et al. 2007a; Nilsen et al. 2008). It was also shown in trout branchial epithelial cells that cortisol increases *cyp11b2* expression (Kolosov and Kelly 2019). These findings suggest the operation of a tissue-level mechanism to regulate cortisol signaling. A better understanding of which branchial cell-types specifically express *cyp11b2* is needed to assess its role in tuning the actions of cortisol on ionocytes.

The role of corticosteroids in regulating permeability of the branchial epithelium has also received considerable attention. This work has largely focused on the FW-adaptive, rather than the SW-adaptive, roles of corticosteroids, as the increased expression of tight-junction proteins generally promotes epithelial tightening. However, “leaky” tight-junction complexes composed of Cldn10s contribute to SW-adaptation by facilitating the paracellular excretion of Na<sup>+</sup> (Tipsmark et al. 2008b; Bui and Kelly 2014). Acclimation to SW increases the expression of *cldn10* isoforms in puffer fish (*Tetraodon nigroviridis*) (Bui et al. 2010) and exposing gill explants to cortisol stimulates multiple *cldn10s* in medaka (Bossus et al. 2017). Cortisol and 11-deoxycorticosterone generally upregulate the expression of Cldns through processes mediated by both the Gr and Mr (Tipsmark et al. 2009; Bui et al. 2010; Chasiotis and Kelly 2011, 2012; Kelly and Chasiotis 2011; Bossus et al. 2017; Kolosov et al. 2017b; Kolosov and Kelly 2019). In sea lamprey, multiple claudins have been identified that are expressed in the gill, and among those investigated, *cldn3* and *-10* orthologs increase their expression after exposure to ion-poor water and exhibit decreases during SW acclimation (Kolosov et al. 2017a, 2020). Future studies in lamprey should seek to address whether 11-deoxycortisol and 11-deoxycorticosterone control branchial barrier functions via Cldns.

Cortisol was the first hormone linked with the expression of branchial Aqps. FW-acclimated eels infused with cortisol show a marked decrease in the expression of *aqp3* in the gill (Cutler et al. 2007) (Fig. 3). Choi et al. (2013) subsequently reported that cortisol diminishes branchial *aqp3* and -8 expression in sockeye salmon (*Oncorhynchus nerka*). These patterns suggest that SW-induced increases in plasma cortisol are responsible for rapidly attenuating *aqp3* expression upon entry into hyperosmotic environments (Cutler and Cramb 2002; Cutler et al. 2007). Furthermore, cortisol blocks the stimulatory action of Prl on *aqp3* (Breves et al. 2016). The regulation of branchial Aqp3 is a clear example of antagonistic, rather than synergistic, roles for cortisol and Prl in promoting salinity acclimation.

### 5.3 Thyroid hormones

In addition to supporting FW acclimation (Section 4.4), there is evidence that thyroid hormones promote SW-adaptive processes by acting directly on ionocytes and through interactions with the Gh/Igf system (McCormick 2001). For example, coho salmon and mummichog increase plasma T<sub>4</sub> levels in response to SW (Knoeppel et al. 1982; Specker and Kobuke 1987), and Atlantic salmon and summer flounder (*Paralichthys dentatus*) treated with T<sub>4</sub> or T<sub>3</sub> exhibit increased SW tolerance (Refstie 1982; Saunders et al. 1985; Schreiber and Specker 1999). Accordingly, when summer flounder and mummichog are treated with thiourea (an inhibitor of T<sub>4</sub> synthesis), they exhibit diminished hyposmoregulatory capacities (Knoeppel et al. 1982; Schreiber and Specker 1999). Thiourea diminishes the SW tolerance of flounder by disrupting the thyroid-mediated development of SW-type ionocytes during metamorphosis (Schreiber and Specker 2000). To our knowledge, there has been no direct assessment of whether the rapid recruitment of SW-type ionocytes that occurs in euryhaline species when they encounter SW is linked with thyroid hormone signaling.

## 6. Future perspectives

The availability of genomic resources and molecular tools over the last two decades has given rise to an increasingly mechanistic understanding of how hormones regulate ionocytes. This trend will undoubtedly continue with manipulative molecular tools such as gene editing ushering in new opportunities to link hormones and their cognate receptors with specific ion transporters. Zebrafish have already proven to be a valuable model for this purpose, supporting progress toward understanding the ontogeny

and function of ion-absorptive ionocytes (Chen et al. 2019). Nonetheless, the poor salinity tolerance of zebrafish precipitates the need for a similarly amenable euryhaline model, a need that Japanese medaka seem poised to fill (Yan and Hwang 2019). In a similar vein, refined methods for primary cell culture of the branchial epithelium would accelerate the use of advanced molecular manipulations; however, progress in this endeavor has been limited.

The various modes by which endocrine factors can affect branchial processes deserve continued attention. For example, it is necessary to better resolve the cytogenic (controlling ionocyte abundance), molecular (controlling the expression of ion transporters), and physiological (controlling the function of ion transporters) actions of hormones (Breves et al 2014a; Shir-Mohammadi and Perry 2020). Important in this endeavor will be the characterization of, 1) the factors influencing the differentiation of SW-type ionocytes from precursor cells (analogous to how Foxi3a and -b regulate FW-type ionocyte differentiation in zebrafish), 2) the regulatory elements in the promoters and distal regulatory regions of genes encoding ion transporters, and 3) the functional elements of the ion transporters themselves (such as the motifs facilitating ATP binding and phosphorylation).

Despite the recent progress, there are still many gaps to fill in the collective understanding of how ionocytes operate – this is especially true for non-teleost fishes. For example, it stands unresolved whether Slc26-family anion exchangers, Clc family Cl<sup>-</sup> channels, and Cftr sustain Cl<sup>-</sup> transport in the ionocytes of lampreys and sturgeons (Ferreira-Martins et al. 2021; Shaughnessy and Breves 2021). We foresee that some of these transporters/channels will emerge as hormone targets. The recent expansion of genomic resources in non-teleosts will certainly support work of this nature (Amemiya et al. 2013; Smith et al. 2013; Braasch et al. 2016; Vialle et al. 2018; Smith et al. 2018; Cheng et al. 2019; Du et al. 2020; Yamaguchi et al. 2020; Marlétaz et al. 2023).

Finally, future work should seek to better understand how systemic hormones interact with the osmotic stress signaling cascades that permit ionocytes to directly perceive salinity changes (Fiol and Kültz 2007). For instance, cortisol promotes the expression of osmotic stress transcription factor 1 (Ostf1) during the acute phase of SW acclimation (McGuire et al. 2010). While Prl inhibits the activity of SW-type ionocytes (Fig. 2), it remains to be seen whether Prl dampens the expression of intracellular and paracrine factors that respond to hyperosmotic conditions (e.g., Ostf1, serum- and glucocorticoid-inducible kinase 1, 14-3-3 proteins, MAPKs, endothelin 1, interleukins,



and tumor necrosis factor  $\alpha$ ) (Fiol and Kültz 2007; Notch et al. 2012; Kültz 2015; Lai et al. 2015). Given the multifactorial nature of osmotic stress signaling (Fiol and Kültz 2007), and the myriad hormones that impact branchial processes (Evans et al. 2005; Takei et al. 2014), it will be interesting to learn the extent to which ionocytes are a hub for interactions between intracellular, paracrine, and systemic signals.

## Competing interests

The authors have no competing interests to declare that are relevant to the content of this article.

## Data availability statement

Data sharing is not applicable to this article as no new data were created or analyzed.

## References

- Agustsson T, Sundell K, Sakamoto T, et al (2001) Growth hormone endocrinology of Atlantic salmon (*Salmo salar*): pituitary gene expression, hormone storage, secretion and plasma levels during parr-smolt transformation. *J Endocrinol* 170:227–234. <https://doi.org/10.1677/joe.0.1700227>
- Amemiya CT, Alföldi J, Lee AP, et al (2013) The African coelacanth genome provides insights into tetrapod evolution. *Nature* 496(7445):311–316. <https://doi.org/10.1038/nature12027>
- Anderson WG (2012) The endocrinology of 1 $\alpha$ -hydroxycorticosterone in elasmobranch fish: a review. *Comp Biochem Physiol A* 162:73–80. <https://doi.org/10.1016/j.cbpa.2011.08.015>
- Aruna A, Nagarajan G, Chang CF (2012a) Involvement of corticotrophin-releasing hormone and corticosteroid receptors in the brain–pituitary–gill of tilapia during the course of seawater acclimation. *J Neuroendocrinol* 24:818–830. <https://doi.org/10.1111/j.1365-2826.2012.02282.x>
- Aruna A, Nagarajan G, Chang CF (2012b) Differential expression patterns and localization of glucocorticoid and mineralocorticoid receptor transcripts in the osmoregulatory organs of tilapia during salinity stress. *Gen Comp Endocrinol* 179:465–476. <https://doi.org/10.1016/j.ygcen.2012.08.028>
- Baker ME (2003) Evolution of glucocorticoid and mineralocorticoid responses: Go fish. *Endocrinology* 144:4223–4225. <https://doi.org/10.1210/en.2003-0843>
- Barany A, Shaughnessy CA, Fuentes J, Mancera JM, McCormick SD. (2020) Osmoregulatory role of the intestine in the sea lamprey (*Petromyzon marinus*). *Am J Physiol Regul Integr Comp Physiol* 318(2):R410–R417. <https://doi.org/10.1152/ajpregu.00033.2019>
- Barany A, Shaughnessy CA, McCormick SD (2021a) Corticosteroid control of Na<sup>+</sup>/K<sup>+</sup>-ATPase in the intestine of the sea lamprey (*Petromyzon marinus*). *Gen Comp Endocrinol* 307:113756. <https://doi.org/10.1016/j.ygcen.2021.113756>
- Barany A, Shaughnessy CA, Pelis RM, et al (2021b) Tissue and salinity specific Na<sup>+</sup>/Cl<sup>-</sup> cotransporter (NCC) orthologues involved in the adaptive osmoregulation of sea lamprey (*Petromyzon marinus*). *Sci Rep* 11:22698. <https://doi.org/10.1038/s41598-021-02125-1>
- Bartels H, Potter IC (2004) Cellular composition and ultrastructure of the gill epithelium of larval and adult lampreys: Implications for osmoregulation in fresh and seawater. *J Exp Biol* 207:3447–3462. <https://doi.org/10.1242/jeb.01157>
- Berishvili G, D'Cotta H, Baroiller JF, Segner H, Reinecke M. (2006) Differential expression of IGF-I mRNA and peptide in the male and female gonad during early development of a bony fish, the tilapia *Oreochromis niloticus*. *Gen Comp Endocrinol* 146(3):204–10. <https://doi.org/10.1016/j.ygcen.2005.11.008>

- Bernard B, Leguen I, Mandiki SNM, et al (2020) Impact of temperature shift on gill physiology during smoltification of Atlantic salmon smolts (*Salmo salar* L.). *Comp Biochem Physiol A Mol Integr Physiol* 244:110685. <https://doi.org/10.1016/j.cbpa.2020.110685>
- Bisbal GA, Specker JL (1991) Cortisol stimulates hypo-osmoregulatory ability in Atlantic salmon, *Salmo salar* L. *J Fish Biol* 39:421–432. <https://doi.org/10.1111/j.1095-8649.1991.tb04373.x>
- Björnsson BT (1997) The biology of salmon growth hormone: from daylight to dominance. *Fish Physiol Biochem* 17:9–24. <https://doi.org/10.1023/A:1007712413908>
- Boeuf G, Le Bail PY, Prunet P (1989) Growth hormone and thyroid hormones during Atlantic salmon, *Salmo salar* L., smolting, and after transfer to seawater. *Aquaculture* 82:257–268. [https://doi.org/10.1016/0044-8486\(89\)90413-4](https://doi.org/10.1016/0044-8486(89)90413-4)
- Bollinger RJ, Ellis LV, Bossus MC, Tipsmark CK (2018) Prolactin controls Na<sup>+</sup>, Cl<sup>-</sup> cotransporter via Stat5 pathway in the teleost gill. *Mol Cell Endocrinol* 477:163–171. <https://doi.org/10.1016/j.mce.2018.06.014>
- Bossus MC, Madsen SS, Tipsmark CK (2015) Functional dynamics of claudin expression in Japanese medaka (*Oryzias latipes*): Response to environmental salinity. *Comp Biochem Physiol A Mol Integr Physiol* 187:74–85. <https://doi.org/10.1016/j.cbpa.2015.04.017>
- Bossus MC, Bollinger RJ, Reed PJ, Tipsmark CK (2017) Prolactin and cortisol regulate branchial claudin expression in Japanese medaka. *Gen Comp Endocrinol* 240:77–83. <https://doi.org/10.1016/j.ygcen.2016.09.010>
- Boyle D, Clifford AM, Orr E, Chamot D, Goss GG (2014) Mechanisms of Cl<sup>-</sup> uptake in rainbow trout: cloning and expression of slc26a6, a prospective Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchanger. *Comp Biochem Physiol A Mol Integr Physiol* 180:43–50. <https://doi.org/10.1016/j.cbpa.2014.11.001>
- Braasch I, Gehrke AR, Smith JJ, et al (2016) The spotted gar genome illuminates vertebrate evolution and facilitates human-teleost comparisons. *Nat Genet* 48(4):427–37. <https://doi.org/10.1038/ng.3526>
- Breves JP (2019) Prolactin controls branchial *clcn2c* but not *atp1a1a.2* in zebrafish *Danio rerio*. *J Fish Biol* 94(1):168–172. <https://doi.org/10.1111/jfb.13854>
- Breves JP, Fox BK, Pierce AL, et al (2010a) Gene expression of growth hormone family and glucocorticoid receptors, osmosensors, and ion transporters in the gill during seawater acclimation of Mozambique tilapia, *Oreochromis mossambicus*. *J Exp Zool* 313A:432–441. <https://doi.org/10.1002/jez.613>
- Breves JP, Hasegawa S, Yoshioka M, et al (2010b) Acute salinity challenges in Mozambique and Nile tilapia: differential responses of plasma prolactin, growth hormone and branchial expression of ion transporters. *Gen Comp Endocrinol* 167:135–142. <https://doi.org/10.1016/j.ygcen.2010.01.022>
- Breves JP, Watanabe S, Kaneko T, et al (2010c) Prolactin restores branchial mitochondrion-rich cells expressing Na<sup>+</sup>/Cl<sup>-</sup> cotransporter in hypophysectomized Mozambique tilapia. *Am J Physiol Regul Integr Comp Physiol* 299(2):R702–10. <https://doi.org/10.1152/ajpregu.00213.2010>
- Breves JP, Seale AP, Helms RE, et al (2011) Dynamic gene expression of GH/PRL-family hormone receptors in gill and kidney during freshwater-acclimation of Mozambique tilapia. *Comp Biochem Physiol A* 158(2):194–200. <https://doi.org/10.1016/j.cbpa.2010.10.030>
- Breves JP, Serizier SB, Goffin V, et al (2013) Prolactin regulates transcription of the ion uptake Na<sup>+</sup>/Cl<sup>-</sup> cotransporter (*ncc*) gene in zebrafish gill. *Mol Cell Endocrinol* 369(1–2):98–106. <https://doi.org/10.1016/j.mce.2013.01.021>
- Breves JP, McCormick SD, Karlstrom RO (2014a) Prolactin and teleost ionocytes: new insights into cellular and molecular targets of prolactin in vertebrate epithelia. *Gen Comp Endocrinol* 203:21–28. <https://doi.org/10.1016/j.ygcen.2013.12.014>
- Breves JP, Seale AP, Moorman BP, et al (2014b) Pituitary control of branchial NCC, NKCC and Na<sup>+</sup>, K<sup>+</sup>-ATPase  $\alpha$ -subunit gene expression in Nile tilapia, *Oreochromis niloticus*. *J Comp Physiol B* 184:513–523. <https://doi.org/10.1007/s00360-014-0817-0>
- Breves JP, Inokuchi M, Yamaguchi Y, et al (2016) Hormonal regulation of aquaporin 3: opposing actions of prolactin and cortisol in tilapia gill. *J Endocrinol* 230(3):325–37. <https://doi.org/10.1530/JOE-16-0162>
- Breves JP, Fujimoto CK, Phipps-Costin SK, et al (2017a) Variation in branchial expression among *insulin-like growth-factor binding proteins (igfbps)* during Atlantic salmon smoltification and seawater exposure. *BMC Physiol* 17:1–11. <https://doi.org/10.1186/s12899-017-0028-5>
- Breves JP, Keith PLK, Hunt BL, et al (2017b) *clc-2c* is regulated by salinity, prolactin and extracellular osmolality in tilapia gill. *J Mol Endocrinol* 59(4):391–402. <https://doi.org/10.1530/JME-17-0144>

- Breves JP, Springer-Miller RH, Chenoweth DA, et al (2020) Cortisol regulates *insulin-like growth-factor binding protein (igfbp)* gene expression in Atlantic salmon parr. *Mol Cell Endocrinol* 518:110989. <https://doi.org/10.1016/j.mce.2020.110989>
- Breves JP, Nelson NN, Koltzenyuk V, et al (2021) Enhanced expression of *ncc1* and *clc2c* in the kidney and urinary bladder accompanies freshwater acclimation in Mozambique tilapia. *Comp Biochem Physiol A* 260:111021. <https://doi.org/10.1016/j.cbpa.2021.111021>
- Breves JP, McKay IS, Koltzenyuk V, et al (2022)  $\text{Na}^+/\text{HCO}_3^-$  cotransporter 1 (*nbce1*) isoform gene expression during smoltification and seawater acclimation of Atlantic salmon. *J Comp Physiol B* 192(5):577-592. <https://doi.org/10.1007/s00360-022-01443-8>
- Breves JP, Runiewicz ER, Richardson SG, et al (2024) Transcriptional regulation of esophageal, intestinal, and branchial solute transporters by salinity, growth hormone, and cortisol in Atlantic salmon. *J Exp Zool Part Ecol Integr Physiol* 341:107–117. <https://doi.org/10.1002/jez.2766>
- Bridgham JT, Carroll SM, Thornton JW (2006) Evolution of hormone-receptor complexity by molecular exploitation. *Science* 312:97–101. <https://doi.org/10.1126/science.1123348>
- Brunelli E, Mauceri A, Salvatore F, et al (2010) Localization of aquaporin 1 and 3 in the gills of the rainbow wrasse *Coris julis*. *Acta Histochem* 112(3):251-8. <https://doi.org/10.1016/j.acthis.2008.11.030>
- Bui P, Bagherie-Lachidan M, Kelly SP (2010) Cortisol differentially alters claudin isoforms in cultured puffer fish gill epithelia. *Mol Cell Endocrinol* 317:120–126. <https://doi.org/10.1016/j.mce.2009.12.002>
- Bui P, Kelly SP (2014) Claudin-6, -10d and -10e contribute to seawater acclimation in the euryhaline puffer fish *Tetraodon nigroviridis*. *J Exp Biol* 217(Pt 10):1758-1767. <https://doi.org/10.1242/jeb.099200>
- Burden CE (1956) The failure of hypophysectomized *Fundulus heteroclitus* to survive in fresh water. *Biol Bull* 110:8-28.
- Bury NR, Sturm A, Le Rouzic P, et al (2003) Evidence for two distinct functional glucocorticoid receptors in teleost fish. *J Mol Endocrinol* 31:141–156. <https://doi.org/10.1677/jme.0.0310141>
- Bystriansky JS, Richards JG, Schulte PM, Ballantyne JS (2006) Reciprocal expression of gill  $\text{Na}^+/\text{K}^+$ -ATPase alpha-subunit isoforms alpha1a and alpha1b during seawater acclimation of three salmonid fishes that vary in their salinity tolerance. *J Exp Biol* 209(Pt 10):1848-1858. <https://doi.org/10.1242/jeb.02188>
- Cerdà J, Finn RN (2010) Piscine aquaporins: an overview of recent advances. *J Exp Zool A* 313(10):623-650. <https://doi.org/10.1002/jez.634>
- Chakraborti P, Mukherjee D (1995) Effects of prolactin and fish pituitary extract on plasma calcium levels in common carp, *Cyprinus carpio*. *Gen Comp Endocrinol* 97(3):320-326. <https://doi.org/10.1006/gcen.1995.1032>
- Chasiotis H, Kelly SP (2011) Effect of cortisol on permeability and tight junction protein transcript abundance in primary cultured gill epithelia from stenohaline goldfish and euryhaline trout. *Gen Comp Endocrinol* 172:494–504. <https://doi.org/10.1016/j.ygcen.2011.04.023>
- Chasiotis H, Kelly SP (2012) Effects of elevated circulating cortisol levels on hydromineral status and gill tight junction protein abundance in the stenohaline goldfish. *Gen Comp Endocrinol* 175:277–283. <https://doi.org/10.1016/j.ygcen.2011.11.024>
- Chasiotis H, Effendi JC, Kelly SP (2009) Occludin expression in goldfish held in ion-poor water. *J Comp Physiol B* 179(2):145-54. <https://doi.org/10.1007/s00360-008-0297-1>
- Chasiotis H, Kolosov D, Bui P, Kelly SP (2012) Tight junctions, tight junction proteins and paracellular permeability across the gill epithelium of fishes: a review. *Respir Physiol Neurobiol* 184:269–281. <https://doi.org/10.1016/j.resp.2012.05.020>
- Chen M, Huang X, Yuen DS, Cheng CH (2011) A study on the functional interaction between the GH/PRL family of polypeptides with their receptors in zebrafish: evidence against GHR1 being the receptor for somatolactin. *Mol Cell Endocrinol* 337(1-2):114-121. <https://doi.org/10.1016/j.mce.2011.02.006>
- Chen YC, Liao BK, Lu YF, et al (2019) Zebrafish *Klf4* maintains the ionocyte progenitor population by regulating epidermal stem cell proliferation and lateral inhibition. *PLoS Genet* 15(4):e1008058. <https://doi.org/10.1371/journal.pgen.1008058>
- Cheng P, Huang Y, Du H, et al (2019) Draft genome and complete *Hox*-cluster characterization of the sterlet (*Acipenser ruthenus*). *Front Genet* 10:776. <https://doi.org/10.3389/fgene.2019.00776>

- Choi YJ, Shin HS, Kim NN, et al (2013) Expression of aquaporin-3 and -8 mRNAs in the parr and smolt stages of sockeye salmon, *Oncorhynchus nerka*: effects of cortisol treatment and seawater acclimation. *Comp Biochem Physiol A* 165(2):228-36. <https://doi.org/10.1016/j.cbpa.2013.03.013>
- Clifford AM, Tresguerres M, Goss GG, Wood CM (2022) A novel K<sup>+</sup>-dependent Na<sup>+</sup> uptake mechanism during low pH exposure in adult zebrafish (*Danio rerio*): new tricks for old dogma. *Acta Physiol (Oxf)* 234(3):e13777. <https://doi.org/10.1111/apha.13777>
- Close DA, Yun S-S, McCormick SD, et al (2010) 11-Deoxycortisol is a corticosteroid hormone in the lamprey. *Proc Natl Acad Sci U S A* 107:13942–13947. <https://doi.org/10.1073/pnas.0914026107>
- Colombe L, Fostier A, Bury N, et al (2000) A mineralocorticoid-like receptor in the rainbow trout, *Oncorhynchus mykiss*: cloning and characterization of its steroid binding domain. *Steroids* 65:319–328. [https://doi.org/10.1016/S0039-128X\(00\)00090-8](https://doi.org/10.1016/S0039-128X(00)00090-8)
- Cruz SA, Chao PL, Hwang PP (2013a) Cortisol promotes differentiation of epidermal ionocytes through Foxi3 transcription factors in zebrafish (*Danio rerio*). *Comp Biochem Physiol A* 164(1):249-257. <https://doi.org/10.1016/j.cbpa.2012.09.011>
- Cruz SA, Lin CH, Chao PL, Hwang PP (2013b) Glucocorticoid receptor, but not mineralocorticoid receptor, mediates cortisol regulation of epidermal ionocyte development and ion transport in zebrafish (*Danio rerio*). *PLoS ONE* 8:e77997. <https://doi.org/10.1371/journal.pone.0077997>
- Cui G, Hong J, Chung-Davidson YW, et al (2019) An ancient CFTR ortholog informs molecular evolution in ABC transporters. *Dev Cell* 51(4):421-430.e3. <https://doi.org/10.1016/j.devcel.2019.09.017>
- Cutler CP, Cramb G (2002) Branchial expression of an aquaporin 3 (AQP-3) homologue is downregulated in the European eel *Anguilla anguilla* following seawater acclimation. *J Exp Biol* 205(Pt 17):2643-51. <https://doi.org/10.1242/jeb.205.17.2643>
- Cutler CP, Phillips C, Hazon N, Cramb G (2007) Cortisol regulates eel (*Anguilla anguilla*) aquaporin 3 (AQP3) mRNA expression levels in gill. *Gen Comp Endocrinol* 152:310-313. <https://doi.org/10.1016/j.ygcen.2007.01.031>
- Dalziel AC, Bittman J, Mandic M, et al (2014) Origins and functional diversification of salinity-responsive Na<sup>+</sup>, K<sup>+</sup> ATPase  $\alpha$ 1 paralogs in salmonids. *Mol Ecol* 23(14):3483-3503. <https://doi.org/10.1111/mec.12828>
- Dauder S, Young G, Hass L, Bern HA (1990) Prolactin receptors in liver, kidney, and gill of the tilapia (*Oreochromis mossambicus*): characterization and effect of salinity on specific binding of iodinated ovine prolactin. *Gen Comp Endocrinol* 77(3):368-377. [https://doi.org/10.1016/0016-6480\(90\)90226-c](https://doi.org/10.1016/0016-6480(90)90226-c)
- Deane EE, Woo NYS (2009) Modulation of fish growth hormone levels by salinity, temperature, pollutants and aquaculture related stress: a review. *Rev Fish Biol Fish* 19:97–120. <https://doi.org/10.1007/s11160-008-9091-0>
- Du K, Stöck M, Kneitz S, et al (2020) The sterlet sturgeon genome sequence and the mechanisms of segmental rediploidization. *Nat Ecol Evol* 4(6):841-852. <https://doi.org/10.1038/s41559-020-1166-x>
- Dymowska AK, Hwang PP, Goss GG (2012) Structure and function of ionocytes in the freshwater fish gill. *Respir Physiol Neurobiol* 184(3):282-292. <https://doi.org/10.1016/j.resp.2012.08.025>
- Eckert SM, Yada T, Shepherd BS, et al (2001) Hormonal control of osmoregulation in the channel catfish *Ictalurus punctatus*. *Gen Comp Endocrinol* 122(3):270-86. <https://doi.org/10.1006/gcen.2001.7633>
- Einarsdóttir IE, Gong N, Jönsson E, et al (2014) Plasma growth hormone-binding protein levels in Atlantic salmon *Salmo salar* during smoltification and seawater transfer. *J Fish Biol* 85:1279–1296. <https://doi.org/10.1111/jfb.12473>
- Ellis LV, Bollinger RJ, Weber HM, et al (2019) Differential expression and localization of branchial AQP1 and AQP3 in Japanese medaka (*Oryzias latipes*). *Cells* 8(5):422. <https://doi.org/10.3390/cells8050422>
- Evans DH, Piermarini PM, Choe KP (2005) The multifunctional fish gill: dominant site of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. *Physiol Rev* 85(1):97-177. <https://doi.org/10.1152/physrev.00050.2003>
- Fargher RC, McKeown BA (1989) The effect of prolactin on calcium homeostasis in coho salmon (*Oncorhynchus kisutch*). *Gen Comp Endocrinol* 73(3):398-403. [https://doi.org/10.1016/0016-6480\(89\)90197-4](https://doi.org/10.1016/0016-6480(89)90197-4)
- Faught E, Vijayan MM (2018) The mineralocorticoid receptor is essential for stress axis regulation in zebrafish larvae. *Sci Rep* 8:18081. <https://doi.org/10.1038/s41598-018-36681-w>
- Ferreira-Martins D, Coimbra J, Antunes C, Wilson JM (2016) Effects of salinity on upstream-migrating, spawning sea lamprey, *Petromyzon marinus*. *Conserv Physiol* 4:1–16. <https://doi.org/10.1093/conphys/cov064>

- Ferreira-Martins D, Wilson JM, Kelly SP, et al (2021) A review of osmoregulation in lamprey. *J Gt Lakes Res* 47:S59–S71. <https://doi.org/10.1016/j.jglr.2021.05.003>
- Ferreira-Martins D, Walton E, Karlstrom RO, et al (2023) The GH/IGF axis in the sea lamprey during metamorphosis and seawater acclimation. *Mol Cell Endocrinol* 111937. <https://doi.org/10.1016/j.mce.2023.111937>
- Fiol DF, Kültz D (2007) Osmotic stress sensing and signaling in fishes. *FEBS J* 274(22):5790–5798. <https://doi.org/10.1111/j.1742-4658.2007.06099.x>
- Fiol DF, Sanmarti E, Sacchi R, Kültz D (2009) A novel tilapia prolactin receptor is functionally distinct from its paralog. *J Exp Biol* 212(Pt 13):2007–2015. <https://doi.org/10.1242/jeb.025601>
- Flik G, Fenwick JC, Wendelaar Bonga SE (1989) Calcitropic actions of prolactin in freshwater North American eel (*Anguilla rostrata* LeSueur). *Am J Physiol* 257(1 Pt 2):R74–79. <https://doi.org/10.1152/ajpregu.1989.257.1.R74>
- Flik G, Atsma W, Fenwick JC, et al (1993) Homologous recombinant growth hormone and calcium metabolism in the tilapia, *Oreochromis mossambicus*, adapted to fresh water. *J Exp Biol* 185:107–119. <https://doi.org/10.1242/jeb.185.1.107>
- Flik G, Rentier-Delrue F, Wendelaar Bonga SE (1994) Calcitropic effects of recombinant prolactins in *Oreochromis mossambicus*. *Am J Physiol* 266(4 Pt 2):R1302–1308. <https://doi.org/10.1152/ajpregu.1994.266.4.R1302>
- Flik G, Klaren, P.H.M., Schoenmakers, et al (1996) Cellular calcium transport in fish: unique and universal mechanisms. *Physiol Biochem Zool* 69:403–417. <http://doi.org/10.1086/physzool.69.2.30164192>
- Flores AM, Shrimpton MJ (2012) Differential physiological and endocrine responses of rainbow trout, *Oncorhynchus mykiss*, transferred from fresh water to ion-poor or salt water. *Gen Comp Endocrinol* 175:244–250. <https://doi.org/10.1016/j.ygcen.2011.11.002>
- Fontaine M, Hatey J (1954) Sur la teneur en 17-hydroxycorticostéroïdes du plasma de saumon (*Salmo salar* L.). *CR Acad Sci Ser D* 239:319–321.
- Fuentes J, Brinca L, Guerreiro PM, Power DM (2010) PRL and GH synthesis and release from the sea bream (*Sparus auratus* L.) pituitary gland *in vitro* in response to osmotic challenge. *Gen Comp Endocrinol* 168(1):95–102. <https://doi.org/10.1016/j.ygcen.2010.04.005>
- Fuller PJ, Yao YZ, Jin R, et al (2019) Molecular evolution of the switch for progesterone and spironolactone from mineralocorticoid receptor agonist to antagonist. *Proc Natl Acad Sci* 201903172. <https://doi.org/10.1073/pnas.1903172116>
- Geering K (2008) Functional roles of Na,K-ATPase subunits. *Curr Opin Nephrol Hypertens* 17(5):526–532. <https://doi.org/10.1097/MNH.0b013e3283036cbf>
- Gilmour KM, Collier CL, Dey CJ, Perry SF (2011) Roles of cortisol and carbonic anhydrase in acid-base compensation in rainbow trout, *Oncorhynchus mykiss*. *J Comp Physiol B* 181(4):501–515. <https://doi.org/10.1007/s00360-010-0540-4>
- Gong N, Ferreira-Martins D, McCormick SD, Sheridan MA (2020) Divergent genes encoding the putative receptors for growth hormone and prolactin in sea lamprey display distinct patterns of expression. *Sci Rep* 10:1674. <https://doi.org/10.1038/s41598-020-58344-5>
- Gong N, Ferreira-Martins D, Norstog JL, et al (2022) Discovery of prolactin-like in lamprey: role in osmoregulation and new insight into the evolution of the growth hormone/prolactin family. *Proc Natl Acad Sci USA* 119:e2212196119. <https://doi.org/10.1073/pnas.2212196119>
- Gray ES, Young G, Bern HA (1990) Radioreceptor assay for growth hormone in coho salmon (*Oncorhynchus kisutch*) and its application to the study of stunting. *J Exp Zool* 256:290–296. <https://doi.org/10.1002/jez.1402560308>
- Guh YJ, Lin CH, Hwang PP (2015) Osmoregulation in zebrafish: ion transport mechanisms and functional regulation. *EXCLI J* 14:627–659. <https://doi.org/10.17179/excli2015-246>
- Herndon TM, McCormick SD, Bern HA (1991) Effects of prolactin on chloride cells in opercular membrane of seawater-adapted tilapia. *Gen Comp Endocrinol* 83(2):283–289. [https://doi.org/10.1016/0016-6480\(91\)90032-2](https://doi.org/10.1016/0016-6480(91)90032-2)
- Hirano T (1986) The spectrum of prolactin action in teleosts. *Prog Clin Biol Res* 205:53–74.
- Hirata T, Kaneko T, Ono T, et al (2003) Mechanism of acid adaptation of a fish living in a pH 3.5 lake. *Am J Physiol Regul Integr Comp Physiol* 284(5):R1199–1212. <https://doi.org/10.1152/ajpregu.00267.2002>

- Hiroi J, McCormick SD (2012) New insights into gill ionocyte and ion transporter function in euryhaline and diadromous fish. *Respir Physiol Neurobiol* 184(3):257-268. <https://doi.org/10.1016/j.resp.2012.07.019>
- Hiroi J, Yasumasu S, McCormick SD, et al (2008) Evidence for an apical Na-Cl cotransporter involved in ion uptake in a teleost fish. *J Exp Biol* 211(Pt 16):2584-2599. <https://doi.org/10.1242/jeb.018663>
- Hoar WS (1988) The physiology of smolting salmonids. In: Hoar WS, Randall DJ (eds) *Fish Physiology*. Academic Press, New York, pp 275–343.
- Hoshijima K, Hirose S (2007) Expression of endocrine genes in zebrafish larvae in response to environmental salinity. *J Endocrinol* 193(3):481-491. <https://doi.org/10.1677/JOE-07-0003>
- Hsiao CD, You MS, Guh YJ, et al (2007) A positive regulatory loop between foxi3a and foxi3b is essential for specification and differentiation of zebrafish epidermal ionocytes. *PLoS One* 2(3):e302. <https://doi.org/10.1371/journal.pone.0000302>
- Hsu HH, Lin LY, Tseng YC, et al (2014) A new model for fish ion regulation: identification of ionocytes in freshwater- and seawater-acclimated medaka (*Oryzias latipes*). *Cell Tissue Res* 357(1):225-243. <https://doi.org/10.1007/s00441-014-1883-z>
- Huang X, Jiao B, Fung CK, et al (2007) The presence of two distinct prolactin receptors in seabream with different tissue distribution patterns, signal transduction pathways and regulation of gene expression by steroid hormones. *J Endocrinol* 194(2):373-92. <https://doi.org/10.1677/JOE-07-0076>
- Hwang PP, Lin LY (2014) Gill ionic transport, acid-base regulation, and nitrogen excretion. In: Evans DH, Claiborne JB, Currie S (eds) *The physiology of fishes*, 4<sup>th</sup> edn. CRC Press, Boca Raton, pp 453
- Inokuchi M, Hiroi J, Watanabe S, et al (2008) Gene expression and morphological localization of NHE3, NCC and NKCC1a in branchial mitochondria-rich cells of Mozambique tilapia (*Oreochromis mossambicus*) acclimated to a wide range of salinities. *Comp Biochem Physiol A* 151(2):151-158. <https://doi.org/10.1016/j.cbpa.2008.06.012>
- Inokuchi M, Hiroi J, Watanabe S, et al (2009) Morphological and functional classification of ion-absorbing mitochondria-rich cells in the gills of Mozambique tilapia. *J Exp Biol* 2212(Pt 7):1003-1010. <https://doi.org/10.1242/jeb.025957>
- Inokuchi M, Breves JP, Moriyama S, et al (2015) Prolactin 177, prolactin 188, and extracellular osmolality independently regulate the gene expression of ion transport effectors in gill of Mozambique tilapia. *Am J Physiol Regul Integr Comp Physiol* 309(10):R1251-1263. <https://doi.org/10.1152/ajpregu.00168.2015>
- Inokuchi M, Hiroi J, Kaneko T (2022) Why can Mozambique tilapia acclimate to both freshwater and seawater? Insights from the plasticity of ionocyte functions in the euryhaline teleost. *Front Physiol* 13:914277. <https://doi.org/10.3389/fphys.2022.914277>
- Ivanis G, Esbaugh AJ, Perry SF (2008) Branchial expression and localization of SLC9A2 and SLC9A3 sodium/hydrogen exchangers and their possible role in acid-base regulation in freshwater rainbow trout (*Oncorhynchus mykiss*). *J Exp Biol* 211(Pt 15):2467-2477. <https://doi.org/10.1242/jeb.017491>
- Jung D, MacIver B, Jackson BP, et al (2012) A novel aquaporin 3 in killifish (*Fundulus heteroclitus*) is not an arsenic channel. *Toxicol Sci* 127(1):101-109. <https://doi.org/10.1093/toxsci/kfs078>
- Kakizawa S, Kaneko T, Hasegawa S, Hirano T (1993) Activation of somatolactin cells in the pituitary of the rainbow trout *Oncorhynchus mykiss* by low environmental calcium. *Gen Comp Endocrinol* 91(3):298-306. <https://doi.org/10.1006/gcen.1993.1130>
- Kaneko T, Hirano T (1983) Role of prolactin and somatolactin in calcium regulation in fish. *J Exp Biol* 184(1):31–45. <https://doi.org/10.1242/jeb.184.1.31>
- Kaneko T, Watanabe S, Lee KM (2008) Functional morphology of mitochondrion-rich cells in euryhaline and stenohaline teleosts. *Aqua-BioSci Monogr* 1:1-62. <https://doi.org/10.5047/absm.2008.00101.0001>
- Katsu Y, Oka K, Baker ME (2018) Evolution of human, chicken, alligator, frog, and zebrafish mineralocorticoid receptors: allosteric influence on steroid specificity. *Sci Signal* 11:35–37. <https://doi.org/10.1126/scisignal.aao1520>
- Kawauchi H, Suzuki K, Yamazaki T, et al (2002) Identification of growth hormone in the sea lamprey, an extant representative of a group of the most ancient vertebrates. *Endocrinology* 143(12):4916-21. <https://doi.org/10.1210/en.2002-220810>
- Kelly SP, Chasiotis H (2011) Glucocorticoid and mineralocorticoid receptors regulate paracellular permeability in a primary cultured gill epithelium. *J Exp Biol* 214:2308–2318. <https://doi.org/10.1242/jeb.055962>

- Kelly SP, Wood CM (2002) Cultured gill epithelia from freshwater tilapia (*Oreochromis niloticus*): effect of cortisol and homologous serum supplements from stressed and unstressed fish. *J Membr Biol* 190(1):29-42. <https://doi.org/10.1007/s00232-002-1020-x>
- Kelly SP, Chow IN, Woo NY (1999) Effects of prolactin and growth hormone on strategies of hypoosmotic adaptation in a marine teleost, *Sparus sarba*. *Gen Comp Endocrinol* 113(1):9-22. <https://doi.org/10.1006/gcen>
- Khodabandeh S, Mosafer S, Khoshnood Z (2009) Effects of cortisol and salinity acclimation on Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> cotransporter gene expression and Na<sup>+</sup>, K<sup>+</sup>-ATPase activity in the gill of Persian sturgeon, *Acipenser persicus*, fry. *Sci Mar* 73S1:111–116. <https://doi.org/10.3989/scimar.2009.73s1111>
- Kiilerich P, Kristiansen K, Madsen SS (2007a) Hormone receptors in gills of smolting Atlantic salmon, *Salmo salar*: expression of growth hormone, prolactin, mineralocorticoid and glucocorticoid receptors and 11 $\beta$ -hydroxysteroid dehydrogenase type 2. *Gen Comp Endocrinol* 152:295–303. <https://doi.org/10.1016/j.ygcen.2006.12.018>
- Kiilerich P, Kristiansen K, Madsen SS (2007b) Cortisol regulation of ion transporter mRNA in Atlantic salmon gill and the effect of salinity on the signaling pathway. *J Endocrinol* 194:417–427. <https://doi.org/10.1677/JOE-07-0185>
- Kiilerich P, Milla S, Sturm A, et al (2011a) Implication of the mineralocorticoid axis in rainbow trout osmoregulation during salinity acclimation. *J Endocrinol* 209:221–235. <https://doi.org/10.1530/JOE-10-0371>
- Kiilerich P, Pedersen SH, Kristiansen K, Madsen SS (2011b) Corticosteroid regulation of Na<sup>+</sup>,K<sup>+</sup>-ATPase  $\alpha$ 1-isoform expression in Atlantic salmon gill during smolt development. *Gen Comp Endocrinol* 170:283–289. <https://doi.org/10.1016/j.ygcen.2010.02.014>
- Kiilerich P, Tipsmark CK, Borski RJ, Madsen SS (2011c) Differential effects of cortisol and 11-deoxycorticosterone on ion transport protein mRNA levels in gills of two euryhaline teleosts, Mozambique tilapia (*Oreochromis mossambicus*) and striped bass (*Morone saxatilis*). *J Endocrinol* 209:115–126. <https://doi.org/10.1530/JOE-10-0326>
- Klaren PH, Guzmán JM, Reutelingsperger SJ, et al (2007) Low salinity acclimation and thyroid hormone metabolizing enzymes in gilthead seabream (*Sparus auratus*). *Gen Comp Endocrinol* 152(2-3):215-22. <https://doi.org/10.1016/j.ygcen.2007.02.010>
- Knoeppel SJ, Atkins DL, Packer RK (1982) The role of the thyroid in osmotic and ionic regulation in *Fundulus heteroclitus* acclimated to freshwater and seawater. *Comp Biochem Physiol A* 73:25-29. [https://doi.org/10.1016/0300-9629\(82\)90087-1](https://doi.org/10.1016/0300-9629(82)90087-1)
- Kolosov D, Bui P, Donini A, et al (2017a) A role for tight junction-associated MARVEL proteins in larval sea lamprey (*Petromyzon marinus*) osmoregulation. *J Exp Biol* 220:3657–3670. <https://doi.org/10.1242/jeb.161562>
- Kolosov D, Donini A, Kelly SP (2017b) Claudin-31 contributes to corticosteroid-induced alterations in the barrier properties of the gill epithelium. *Mol Cell Endocrinol* 439:457–466. <https://doi.org/10.1016/j.mce.2016.10.034>
- Kolosov D, Kelly SP (2019) The mineralocorticoid receptor contributes to barrier function of a model fish gill epithelium. *J Exp Biol* 222(Pt 11):jeb192096. <https://doi.org/10.1242/jeb.192096>
- Kolosov D, Bui P, Wilkie MP, Kelly SP (2020) Claudins of sea lamprey (*Petromyzon marinus*) – organ-specific expression and transcriptional responses to water of varying ion content. *J Fish Biol* 768–781. <https://doi.org/10.1111/jfb.14274>
- Kolosov D, Kelly SP (2017) Claudin-8d is a cortisol-responsive barrier protein in the gill epithelium of trout. *J Mol Endocrinol* 59(3):299-310. <https://doi.org/10.1530/JME-17-0108>
- Kültz D (2015) Physiological mechanisms used by fish to cope with salinity stress. *J Exp Biol* 218(Pt 12):1907-1914. <https://doi.org/10.1242/jeb.118695>
- Kumai Y, Bahubeshi A, Steele S, Perry SF (2011) Strategies for maintaining Na<sup>+</sup> balance in zebrafish (*Danio rerio*) during prolonged exposure to acidic water. *Comp Biochem Physiol A* 160(1):52-62. <https://doi.org/10.1016/j.cbpa.2011.05.001>
- Kumai Y, Nesan D, Vijayan MM, Perry SF (2012) Cortisol regulates Na<sup>+</sup> uptake in zebrafish, *Danio rerio*, larvae via the glucocorticoid receptor. *Mol Cell Endocrinol* 364(1-2):113-25. <https://doi.org/10.1016/j.mce.2012.08.017>
- Lai KP, Li JW, Gu J, et al (2015) Transcriptomic analysis reveals specific osmoregulatory adaptive responses in gill mitochondria-rich cells and pavement cells of the Japanese eel. *BMC Genomics* 16:1072. <https://doi.org/10.1186/s12864-015-2271-0>
- Langhorn P, Simpson TH (1986) The interrelationship of cortisol, gill Na<sup>+</sup>/K<sup>+</sup> ATPase, and homeostasis during the parr-smolt transformation of Atlantic salmon (*Salmo salar* L.). *Gen Comp Endocrinol* 61:203–213. [https://doi.org/10.1016/0016-6480\(86\)90198-X](https://doi.org/10.1016/0016-6480(86)90198-X)

- Lee KM, Kaneko T, Aida K (2006) Prolactin and prolactin receptor expressions in a marine teleost, pufferfish *Takifugu rubripes*. *Gen Comp Endocrinol* 146(3):318-28. <https://doi.org/10.1016/j.ygcen.2005.12.003>
- Leguen I, Le Cam A, Montfort J, et al (2015) Transcriptomic analysis of trout gill ionocytes in fresh water and sea water using laser capture microdissection combined with microarray analysis. *PLoS One* 10(10):e0139938. <https://doi.org/10.1371/journal.pone.0139938>
- Liao BK, Deng AN, Chen SC, et al (2007) Expression and water calcium dependence of calcium transporter isoforms in zebrafish gill mitochondrion-rich cells. *BMC Genomics* 8:354. <https://doi.org/10.1186/1471-2164-8-354>
- Liao BK, Chen RD, Hwang PP (2009) Expression regulation of Na<sup>+</sup>-K<sup>+</sup>-ATPase alpha1-subunit subtypes in zebrafish gill ionocytes. *Am J Physiol Regul Integr Comp Physiol* 296(6):R1897-906. <https://doi.org/10.1152/ajpregu.00029.2009>
- Lignot JH, Cutler CP, Hazon N, Cramb G (2002) Immunolocalisation of aquaporin 3 in the gill and the gastrointestinal tract of the European eel *Anguilla anguilla* (L.). *J Exp Biol* 205(Pt 17):2653-2663. <https://doi.org/10.1242/jeb.205.17.2653>
- Lin CH, Hwang PP (2016) The control of calcium metabolism in zebrafish (*Danio rerio*). *Int J Mol Sci* 17(11):1783. <https://doi.org/10.3390/ijms17111783>
- Link K, Berishvili G, Shved N, et al (2010) Seawater and freshwater challenges affect the insulin-like growth factors IGF-I and IGF-II in liver and osmoregulatory organs of the tilapia. *Mol Cell Endocrinol* 327(1-2):40-46. <https://doi.org/10.1016/j.mce.2010.05.011>
- Link K, Shved N, Serrano N, et al (2022) Effects of seawater and freshwater challenges on the Gh/Igf system in the saline-tolerant blackchin tilapia (*Sarotherodon melanotheron*). *Front Endocrinol* 13:976488. <https://doi.org/10.3389/fendo.2022.976488>
- López-Bojórquez L, Villalobos P, García-G C, et al (2007) Functional identification of an osmotic response element (ORE) in the promoter region of the killifish deiodinase 2 gene (FhDio2). *J Exp Biol*. 210(Pt 17):3126-3132. <https://doi.org/10.1242/jeb.004150>
- Loretz CA, Bern HA (1982) Prolactin and osmoregulation in vertebrates: an update. *Neuroendocrinology* 35(4):292-304. <https://doi.org/10.1159/000123397>
- Mackie P, Wright PA, Glebe BD, Ballantyne JS (2005) Osmoregulation and gene expression of Na<sup>+</sup>/K<sup>+</sup> ATPase in families of Atlantic salmon (*Salmo salar*) smolts. *Can J Fish Aquat Sci* 62(11):2661-2672. <https://doi.org/10.1139/f05-168>
- Madsen SS (1990a) Enhanced hypoosmoregulatory response to growth hormone after cortisol treatment in immature rainbow trout, *Salmo gairdneri*. *Fish Physiol Biochem* 8:271–279. <https://doi.org/10.1007/BF00003422>
- Madsen SS (1990b) The role of cortisol and growth hormone in seawater adaptation and development of hypoosmoregulatory mechanisms in sea trout parr (*Salmo trutta trutta*). *Gen Comp Endocrinol* 79:1–11. [https://doi.org/10.1016/0016-6480\(90\)90082-W](https://doi.org/10.1016/0016-6480(90)90082-W)
- Madsen SS, Bern HA (1993) In-vitro effects of insulin-like growth factor-I on gill Na<sup>+</sup>,K<sup>+</sup>-ATPase in coho salmon, *Oncorhynchus kisutch*. *J Endocrinol* 138(1):23-30. <https://doi.org/10.1677/joe.0.1380023>
- Madsen SS, Kiilerich P, Tipsmark CK (2009) Multiplicity of expression of Na<sup>+</sup>,K<sup>+</sup>-ATPase alpha-subunit isoforms in the gill of Atlantic salmon (*Salmo salar*): cellular localisation and absolute quantification in response to salinity change. *J Exp Biol* 212(Pt 1):78-88. <https://doi.org/10.1242/jeb.024612>. PMID: 19088213
- Madsen SS, Korsgaard B (1991) Opposite effects of 17β-estradiol and combined growth hormone-cortisol treatment on hypo-osmoregulatory performance in sea trout presmolts, *Salmo trutta*. *Gen Comp Endocrinol* 83:276–282. [https://doi.org/10.1016/0016-6480\(91\)90031-Z](https://doi.org/10.1016/0016-6480(91)90031-Z)
- Mancera JM, Carrión RL, del Río MPM (2002) Osmoregulatory action of PRL, GH, and cortisol in the gilthead seabream (*Sparus aurata* L.). *Gen Comp Endocrinol* 129(2):95-103. [https://doi.org/10.1016/S0016-6480\(02\)00522-1](https://doi.org/10.1016/S0016-6480(02)00522-1)
- Mancera JM, McCormick SD (1998) Evidence for growth hormone/insulin-like growth factor I axis regulation of seawater acclimation in the euryhaline teleost *Fundulus heteroclitus*. *Gen Comp Endocrinol* 111:103–112. <https://doi.org/10.1006/gcen.1998.7086>
- Manzon LA (2002) The role of prolactin in fish osmoregulation: a review. *Gen Comp Endocrinol* 125(2):291-310. <https://doi.org/10.1006/gcen.2001.7746>
- Marlétaz F, de la Calle-Mustienes E, Acemel RD, et al (2023) The little skate genome and the evolutionary emergence of wing-like fins. *Nature* 616(7957):495-503. <https://doi.org/10.1038/s41586-023-05868-1>
- Marsigliante S, Barker S, Jimenez E, Storelli C (2000) Glucocorticoid receptors in the euryhaline teleost *Anguilla anguilla*. *Mol Cell Endocrinol* 162:193–201. [https://doi.org/10.1016/S0303-7207\(99\)00262-2](https://doi.org/10.1016/S0303-7207(99)00262-2)



- Maule AG, Schreck CB (1990) Glucocorticoid receptors in leukocytes and gill of juvenile coho salmon (*Oncorhynchus kisutch*). *Gen Comp Endocrinol* 77:448–455. [https://doi.org/10.1016/0016-6480\(90\)90236-f](https://doi.org/10.1016/0016-6480(90)90236-f)
- Marshall WS, Grosell, M (2006). Ion transport, osmoregulation, and acid-base balance. In: Evans DH & Claiborne JB (eds) *The physiology of fishes*, 3<sup>rd</sup> edn. CRC Press, Boca Raton, pp 177–230
- Marshall WS, Breves JP, Doohan EM, et al (2018) *claudin-10* isoform expression and cation selectivity change with salinity in salt-secreting epithelia of *Fundulus heteroclitus*. *J Exp Biol* 221(Pt 1):jeb168906. <https://doi.org/10.1242/jeb.168906>
- Mazurais D, Ducouret B, Tujague M, et al (1998) Regulation of the glucocorticoid receptor mRNA levels in the gills of Atlantic salmon (*Salmo salar*) during smoltification. *Bull Fr Pêche Piscic* 350-351:499-510. <https://doi.org/10.1051/kmae:1998019>
- McCormick SD (1996) Effects of growth hormone and insulin-like growth factor I on salinity tolerance and gill Na<sup>+</sup>, K<sup>+</sup>-ATPase in Atlantic salmon (*Salmo salar*): interaction with cortisol. *Gen Comp Endocrinol* 101:3–11. <https://doi.org/10.1006/gcen.1996.0002>
- McCormick SD (2001) Endocrine control of osmoregulation in teleost fish. *Am Zool* 41:781–794. <https://doi.org/10.1093/icb/41.4.781>
- McCormick SD (2013) Smolt physiology and endocrinology. In: McCormick SD, Farrell AP, Brauner CJ (eds) *Fish Physiology: Euryhaline Fishes*. Academic Press, Inc., Amsterdam, pp 191–251.
- McCormick SD, Bern HA (1989) *In vitro* stimulation of Na<sup>+</sup>-K<sup>+</sup>-ATPase activity and ouabain binding by cortisol in coho salmon gill. *Am J Physiol - Regul Integr Comp Physiol* 256:R707–715. <https://doi.org/10.1152/ajpregu.1989.256.3.r707>
- McCormick SD, Dickhoff WW, Duston J, et al (1991) Developmental differences in the responsiveness of gill Na<sup>+</sup>, K<sup>+</sup>-ATPase to cortisol in salmonids. *Gen Comp Endocrinol* 84:308–317. [https://doi.org/10.1016/0016-6480\(91\)90054-A](https://doi.org/10.1016/0016-6480(91)90054-A)
- McCormick SD, Regish A, O'Dea MF, Shrimpton JM (2008) Are we missing a mineralocorticoid in teleost fish? Effects of cortisol, deoxycorticosterone and aldosterone on osmoregulation, gill Na<sup>+</sup>,K<sup>+</sup>-ATPase activity and isoform mRNA levels in Atlantic salmon. *Gen Comp Endocrinol* 157:35–40. <https://doi.org/10.1016/j.ygcen.2008.03.024>
- McCormick SD, Regish AM, Christensen AK (2009) Distinct freshwater and seawater isoforms of Na<sup>+</sup>/K<sup>+</sup>-ATPase in gill chloride cells of Atlantic salmon. *J Exp Biol* 212(Pt 24):3994–4001. <https://doi.org/10.1242/jeb.037275>
- McCormick SD, Regish AM, Christensen AK, Björnsson BT (2013) Differential regulation of sodium–potassium pump isoforms during smolt development and seawater exposure of Atlantic salmon. *J Exp Biol* 216:1142–1151. <https://doi.org/10.1242/jeb.080440>
- McCormick SD, Saunders RL (1987) Preparatory physiological adaptations for marine life of salmonids: osmoregulation, growth, and metabolism. *Am Fish Soc Symp* 1:211–229. [https://doi.org/10.1016/0022-4596\(90\)90074-8](https://doi.org/10.1016/0022-4596(90)90074-8)
- McCormick SD, Shrimpton JM, Moriyama S, Björnsson BT (2007) Differential hormonal responses of Atlantic salmon parr and smolt to increased daylength: a possible developmental basis for smolting. *Aquaculture* 273:337–344. <https://doi.org/10.1016/j.aquaculture.2007.10.015>
- McCormick SD, Taylor ML, Regish AM (2020) Cortisol is an osmoregulatory and glucose-regulating hormone in Atlantic sturgeon, a basal ray-finned fish. *J Exp Biol* 223:jeb220251. <https://doi.org/10.1242/jeb.220251>
- McGuire A, Aluru N, Takemura A, et al (2010) Hyperosmotic shock adaptation by cortisol involves upregulation of branchial osmotic stress transcription factor 1 gene expression in Mozambique tilapia. *Gen Comp Endocrinol* 165(2):321–9. <https://doi.org/10.1016/j.ygcen.2009.07.016>
- Mizuno S, Ura K, Onodera Y, et al (2001) Changes in transcript levels of gill cortisol receptor during smoltification in wild masu salmon, *Oncorhynchus masou*. *Zoolog Sci* 18:853–860. <https://doi.org/10.2108/zsj.18.853>
- Motoshima T, Nagashima A, Ota C, et al (2023) Na<sup>+</sup>/Cl<sup>-</sup> cotransporter 2 is not fish-specific and is widely found in amphibians, non-avian reptiles, and select mammals. *Physiol Genomics* 55(3):113–131. <https://doi.org/10.1152/physiolgenomics.00143.2022>
- Moyle PB, Cech JJ (2004) *Fishes: an introduction to ichthyology*, 4<sup>th</sup> edn. Prentice Hall, Upper Saddle River, NJ
- Nilsen TO, Ebbesson LO, Madsen SS, et al (2007) Differential expression of gill Na<sup>+</sup>,K<sup>+</sup>-ATPase alpha- and beta-subunits, Na<sup>+</sup>,K<sup>+</sup>,2Cl<sup>-</sup> cotransporter and CFTR anion channel in juvenile anadromous and landlocked Atlantic salmon *Salmo salar*. *J Exp Biol* 210(Pt 16):2885–96. <https://doi.org/10.1242/jeb.002873>

- Nilsen TO, Ebbesson LOE, Kiilerich P, et al (2008) Endocrine systems in juvenile anadromous and landlocked Atlantic salmon (*Salmo salar*): seasonal development and seawater acclimation. *Gen Comp Endocrinol* 155:762–772. <https://doi.org/10.1016/j.ygcen.2007.08.006>
- Notch EG, Chapline C, Flynn E, et al (2012) Mitogen activated protein kinase 14-1 regulates serum glucocorticoid kinase 1 during seawater acclimation in Atlantic killifish, *Fundulus heteroclitus*. *Comp Biochem Physiol A* 162(4):443–448. <https://doi.org/10.1016/j.cbpa.2012.04.025>
- Nozaki M, Ominato K, Shimotani T, et al (2008) Identity and distribution of immunoreactive adenohypophyseal cells in the pituitary during the life cycle of sea lampreys, *Petromyzon marinus*. *Gen Comp Endocrinol* 155:403–412. <https://doi.org/10.1016/j.ygcen.2007.07.012>
- Pang PK, Schreibman MP, Balbontin F, Pang RK (1978) Prolactin and pituitary control of calcium regulation in the killifish, *Fundulus heteroclitus*. *Gen Comp Endocrinol* 36(2):306–16. [https://doi.org/10.1016/0016-6480\(78\)90037-0](https://doi.org/10.1016/0016-6480(78)90037-0)
- Parks SK, Tresguerres M, Goss GG (2007) Interactions between Na<sup>+</sup> channels and Na<sup>+</sup>-HCO<sub>3</sub><sup>-</sup> cotransporters in the freshwater fish gill MR cell: a model for transepithelial Na<sup>+</sup> uptake. *Am J Physiol Cell Physiol* 292(2):C935–44. <https://doi.org/10.1152/ajpcell.00604.2005>
- Pavlovic D, Fuller W, Shattock MJ (2013) Novel regulation of cardiac Na pump via phospholemman. *J Mol Cell Cardiol* 61:83–93. <https://doi.org/10.1016/j.yjmcc.2013.05.002>
- Pelis RM, McCormick SD (2001) Effects of growth hormone and cortisol on Na<sup>+</sup>–K<sup>+</sup>–2Cl<sup>-</sup> cotransporter localization and abundance in the gills of Atlantic salmon. *Gen Comp Endocrinol* 124:134–143. <https://doi.org/10.1006/gcen.2001.7703>
- Pérez-Rius C, Gaitán-Peñas H, Estévez R, Barrallo-Gimeno A (2015) Identification and characterization of the zebrafish CIC-2 chloride channel orthologs. *Pflugers Arch* 467(8):1769–1781. <https://doi.org/10.1007/s00424-014-1614-z>
- Perry SF, Goss GG, Fenwick JC (1992) Interrelationships between gill chloride cell morphology and calcium uptake in freshwater teleosts. *Fish Physiol Biochem* 10(4):327–337. <https://doi.org/10.1007/BF00004482>
- Peter MCS, Lock RA, Wendelaar Bonga SE (2000) Evidence for an osmoregulatory role of thyroid hormones in the freshwater Mozambique tilapia *Oreochromis mossambicus*. *Gen Comp Endocrinol* 120(2):157–67. <https://doi.org/10.1006/gcen.2000.7542>
- Pickford GE (1953) A study of the hypophysectomized male *Fundulus heteroclitus* (Linn.) *Bull Bingham Oceanogr Collect* 14(2):5–45.
- Pickford GE, Atz JW (1957) The physiology of the pituitary gland of fishes. New York Zoological Society, New York.
- Pickford GE, Griffith RW, Torretti J, et al (1970) Branchial reduction and renal stimulation of (Na<sup>+</sup>, K<sup>+</sup>)-ATPase by prolactin in hypophysectomized killifish in fresh water. *Nature* 228(5269):378–379. <https://doi.org/10.1038/228378a0>
- Pickford GE, Pang PKT, Weinstein E, et al (1970) The response of the hypophysectomized cyprinodont, *Fundulus heteroclitus*, to replacement therapy with cortisol: effects on blood serum and sodium-potassium activated adenosine triphosphatase in the gills, kidney, and intestinal mucosa. *Gen Comp Endocrinol* 14:524–534. [https://doi.org/10.1016/0016-6480\(70\)90036-5](https://doi.org/10.1016/0016-6480(70)90036-5)
- Pickford GE, Phillips JG (1959) Prolactin, a factor in promoting survival of hypophysectomized killifish in fresh water. *Science* 130:454–455. <https://doi.org/10.1126/science.130.3373.454>
- Pierce AL, Fox BK, Davis LK, et al (2007) Prolactin receptor, growth hormone receptor, and putative somatolactin receptor in Mozambique tilapia: tissue specific expression and differential regulation by salinity and fasting. *Gen Comp Endocrinol* 154(1–3):31–40. <https://doi.org/10.1016/j.ygcen.2007.06.023>
- Pierce AL, Breves JP, Moriyama S, et al (2011) Differential regulation of Igf1 and Igf2 mRNA levels in tilapia hepatocytes: effects of insulin and cortisol on GH sensitivity. *J Endocrinol* 211(2):201–210. <https://doi.org/10.1530/JOE-10-0456>
- Pisam M, Auperin B, Prunet P, et al (1993) Effects of prolactin on alpha and beta chloride cells in the gill epithelium of the saltwater adapted tilapia "*Oreochromis niloticus*". *Anat Rec* 235(2):275–84. <https://doi.org/10.1002/ar.1092350211>
- Poppinga J, Kittilson J, McCormick SD, Sheridan MA (2007) Effects of somatostatin on the growth hormone-insulin-like growth factor axis and seawater adaptation of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 273:312–319. <https://doi.org/10.1016/j.aquaculture.2007.10.021>
- Potts WTW, Evans DH (1966). The effects of hypophysectomy and bovine prolactin on salt fluxes in fresh-water-adapted *Fundulus heteroclitus*. *Biol Bull* 131:362–368.

- Prunet P, Auperin B (1994) Prolactin receptors. In: Sherwood NM, Hew CL (eds), Fish physiology, Vol. 13: Molecular Endocrinology of Fish. Academic Press, New York, pp. 367-391
- Prunet P, Boeuf G, Bolton JP, Young G (1989) Smoltification and seawater adaptation in Atlantic salmon (*Salmo salar*): plasma prolactin, growth hormone, and thyroid hormones. Gen Comp Endocrinol 74:355–364. [https://doi.org/10.1016/S0016-6480\(89\)80031-0](https://doi.org/10.1016/S0016-6480(89)80031-0)
- Prunet P, Pisam M, Claireaux JP, et al (1994) Effects of growth hormone on gill chloride cells in juvenile Atlantic salmon (*Salmo salar*). Am J Physiol-Regul Integr Comp Physiol 266:R850–R857. <https://doi.org/10.1152/ajpregu.1994.266.3.R850>
- Prunet P, Sturm A, Milla S (2006) Multiple corticosteroid receptors in fish: from old ideas to new concepts. Gen Comp Endocrinol 147:17–23. <https://doi.org/10.1016/j.ygcen.2006.01.015>
- Rai S, Szeitz A, Roberts BW, et al (2015) A putative corticosteroid hormone in Pacific lamprey, *Entosphenus tridentatus*. Gen Comp Endocrinol 212:178-184. <https://doi.org/10.1016/j.ygcen.2014.06.019>
- Refstie T (1982) The effect of feeding thyroid hormones on saltwater tolerance and growth rate of Atlantic salmon. Can J Zool 60(11):2706-2712. <https://doi.org/10.1139/z82-346>
- Reindl KM, Sheridan MA (2012) Peripheral regulation of the growth hormone-insulin-like growth factor system in fish and other vertebrates. Comp Biochem Physiol A 163(3-4):231-245. <https://doi.org/10.1016/j.cbpa.2012.08.003>
- Reinecke M, Schmid A, Ermatinger R, Löffing-Cueni D (1997) Insulin-like growth factor I in the teleost *Oreochromis mossambicus*, the tilapia: gene sequence, tissue expression, and cellular localization. Endocrinology 138(9):3613-9. <https://doi.org/10.1210/endo.138.9.5375>
- Reis-Santos P, McCormick SD, Wilson JM (2008) Ionoregulatory changes during metamorphosis and salinity exposure of juvenile sea lamprey (*Petromyzon marinus* L.). J Exp Biol 211:978–988. <https://doi.org/10.1242/jeb.014423>
- Ren J, Chung-Davidson YW, Yeh CY, et al (2015) Genome-wide analysis of the ATP-binding cassette (ABC) transporter gene family in sea lamprey and Japanese lamprey. BMC Genomics 16(1):436. <https://doi.org/10.1186/s12864-015-1677-z>
- Rhee JS, Kim RO, Seo JS, et al (2010) Effects of salinity and endocrine-disrupting chemicals on expression of prolactin and prolactin receptor genes in the euryhaline hermaphroditic fish, *Kryptolebias marmoratus*. Comp Biochem Physiol C 152(4):413-23. <https://doi.org/10.1016/j.cbpc.2010.07.001>
- Richards JG, Semple JW, Bystriansky JS, Schulte PM (2003) Na<sup>+</sup>/K<sup>+</sup>-ATPase alpha-isoform switching in gills of rainbow trout (*Oncorhynchus mykiss*) during salinity transfer. J Exp Biol 206(Pt 24):4475-4486. <https://doi.org/10.1242/jeb.00701>
- Rouzic PL, Sandra O, Grosclaude J, et al (2002) Evidence of rainbow trout prolactin interaction with its receptor through unstable homodimerisation. Mol Cell Endocrinol 172(1-2):105-113. [https://doi.org/10.1016/S0303-7207\(00\)00377-4](https://doi.org/10.1016/S0303-7207(00)00377-4)
- Ruhr IM, Wood CM, Schauer KL, et al (2020) Is aquaporin-3 involved in water-permeability changes in the killifish during hypoxia and normoxic recovery, in freshwater or seawater? J Exp Zool A 333(7):511-525. <https://doi.org/10.1002/jez.2393>
- Saito K, Nakamura N, Ito Y, et al (2010) Identification of zebrafish Fxyd11a protein that is highly expressed in ion-transporting epithelium of the gill and skin and its possible role in ion homeostasis. Front Physiol 1:129. <https://doi.org/10.3389/fphys.2010.00129>
- Sakamoto T, Hirano T (1991) Growth hormone receptors in the liver and osmoregulatory organs of rainbow trout: characterization and dynamics during adaptation to seawater. J Endocrinol 130:425–433. <https://doi.org/10.1677/joe.0.1300425>
- Sakamoto T, McCormick SD (2006) Prolactin and growth hormone in fish osmoregulation. Gen Comp Endocrinol 147(1):24-30. <https://doi.org/10.1016/j.ygcen.2005.10.008>
- Sakamoto T, Shepherd BS, Madsen SS, et al (1997) Osmoregulatory actions of growth hormone and prolactin in an advanced teleost. Gen Comp Endocrinol 106(1):95-101. <https://doi.org/10.1006/gcen.1996.6854>
- Santos CR, Ingleton PM, Cavaco JE, et al (2001) Cloning, characterization, and tissue distribution of prolactin receptor in the sea bream (*Sparus aurata*). Gen Comp Endocrinol 121(1):32-47. <https://doi.org/10.1006/gcen.2000.7553>
- Saunders RL, McCormick SD, Henderson EB, et al (1985) The effect of orally administered 3,5,3'-triiodo-L-thyronine on growth and salinity tolerance of Atlantic salmon (*Salmo salar* L.). Aquaculture 45:143-156. [https://doi.org/10.1016/0044-8486\(85\)90265-0](https://doi.org/10.1016/0044-8486(85)90265-0)

- Schultz ET, McCormick SD (2013) Euryhalinity in an evolutionary context. In: McCormick SD, Farrell AP, Brauner CJ (eds) Euryhaline fishes. Elsevier, New York, pp 477-529
- Scott GR, Keir KR, Schulte PM (2005) Effects of spironolactone and RU486 on gene expression and cell proliferation after freshwater transfer in the euryhaline killifish. *J Comp Physiol B* 175:499–510. <https://doi.org/10.1007/s00360-005-0014-2>
- Schreiber AM, Specker JL (1999) Metamorphosis in the summer flounder, *Paralichthys dentatus*: thyroidal status influences salinity tolerance. *J Exp Zool* 284(4):414-424. [https://doi.org/10.1002/\(sici\)1097-010x\(19990901\)284:4<414::aid-jez8>3.0.co;2-e](https://doi.org/10.1002/(sici)1097-010x(19990901)284:4<414::aid-jez8>3.0.co;2-e)
- Schreiber AM, Specker JL (2000) Metamorphosis in the summer flounder, *Paralichthys dentatus*: thyroidal status influences gill mitochondria-rich cells. *Gen Comp Endocrinol* 117(2):238-250. <https://doi.org/10.1006/gcen.1999.7407>
- Seale AP, Riley LG, Leedom TA, et al (2002) Effects of environmental osmolality on release of prolactin, growth hormone and ACTH from the tilapia pituitary. *Gen Comp Endocrinol* 128:91-101. [https://doi.org/10.1016/S0016-6480\(02\)00027-8](https://doi.org/10.1016/S0016-6480(02)00027-8)
- Seale AP, Watanabe S, Grau EG (2012) Osmoreception: perspectives on signal transduction and environmental modulation. *Gen Comp Endocrinol* 176(3):354-360. <https://doi.org/10.1016/j.ygcen.2011.10.005>
- Seale AP, Pavlosky KK, Celino-Brady FT, et al (2019) Systemic versus tissue-level prolactin signaling in a teleost during a tidal cycle. *J Comp Physiol B* 189(5):581-594. <https://doi.org/10.1007/s00360-019-01233-9>
- Seale LA, Gilman CL, Zavacki AM, et al (2021) Regulation of thyroid hormones and branchial iodothyronine deiodinases during freshwater acclimation in tilapia. *Mol Cell Endocrinol* 538:111450. <https://doi.org/10.1016/j.mce.2021.111450>
- Shaughnessy CA, Barany A, McCormick SD (2020) 11-Deoxycortisol controls hydromineral balance in the most basal osmoregulating vertebrate, sea lamprey (*Petromyzon marinus*). *Sci Rep* 10:12148. <https://doi.org/10.1038/s41598-020-69061-4>
- Shaughnessy CA, Breves JP (2021) Molecular mechanisms of Cl<sup>-</sup> transport in fishes: New insights and their evolutionary context. *J Exp Zool A* 335(2):207-216. <https://doi.org/10.1002/jez.2428>
- Shaughnessy CA, McCormick SD (2020) Functional characterization and osmoregulatory role of the Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporter in the gill of sea lamprey (*Petromyzon marinus*), a basal vertebrate. *Am J Physiol Regul Integr Comp Physiol* 318(1):R17-R29. <https://doi.org/10.1152/ajpregu.00125.2019>
- Shimomura T, Nakajima T, Horikoshi M, et al (2012) Relationships between gill Na<sup>+</sup>,K<sup>+</sup>-ATPase activity and endocrine and local insulin-like growth factor-I levels during smoltification of masu salmon (*Oncorhynchus masou*). *Gen Comp Endocrinol* 178:427–435. <https://doi.org/10.1016/j.ygcen.2012.06.011>
- Shir-Mohammadi K, Perry SF (2020) Expression of ion transport genes in ionocytes isolated from larval zebrafish (*Danio rerio*) exposed to acidic or Na<sup>+</sup>-deficient water. *Am J Physiol Regul Integr Comp Physiol* 319(4):R412-R427. <https://doi.org/10.1152/ajpregu.00095.2020>
- Shrimpton JM, Bernier NJ, Randall DJ (1994) Changes in cortisol dynamics in wild and hatchery-reared juvenile coho salmon (*Oncorhynchus kisutch*) during smoltification. *Can J Fish Aquat Sci* 51:2179–2187. <https://doi.org/10.1139/f94-219>
- Shrimpton JM, McCormick SD (1998) Regulation of gill cytosolic corticosteroid receptors in juvenile Atlantic salmon: interaction effects of growth hormone with prolactin and triiodothyronine. *Gen Comp Endocrinol* 112(2):262-274. <https://doi.org/10.1006/gcen.1998.7172>
- Shrimpton J, McCormick SD (1999) Responsiveness of gill Na<sup>+</sup>/K<sup>+</sup>-ATPase to cortisol is related to gill corticosteroid receptor concentration in juvenile rainbow trout. *J Exp Biol* 202:987–995. <https://doi.org/10.1242/jeb.202.8.987>
- Shrimpton JM, Randall DJ (1994) Downregulation of corticosteroid receptors in gills of coho salmon due to stress and cortisol treatment. *Am J Physiol - Regul Integr Comp Physiol* 267:R432-438. <https://doi.org/10.1152/ajpregu.1994.267.2.R432>
- Singer TD, Finstad B, McCormick SD, et al (2003) Interactive effects of cortisol treatment and ambient seawater challenge on gill Na<sup>+</sup>,K<sup>+</sup>-ATPase and CFTR expression in two strains of Atlantic salmon smolts. *Aquaculture* 222:15–28. [https://doi.org/10.1016/S0044-8486\(03\)00099-1](https://doi.org/10.1016/S0044-8486(03)00099-1)
- Sloman KA, Desforges PR, Gilmour KM (2001) Evidence for a mineralocorticoid-like receptor linked to branchial chloride cell proliferation in freshwater rainbow trout. *J Exp Biol* 204:3953–3961. <https://doi.org/10.1242/jeb.204.22.3953>
- Smith JJ, Kuraku S, Holt C, et al (2013) Sequencing of the sea lamprey (*Petromyzon marinus*) genome provides insights into vertebrate evolution. *Nat Genet* 45(4):415-421. <https://doi.org/10.1038/ng.2568>

- Smith JJ, Timoshevskaya N, Ye C, et al (2018) The sea lamprey germline genome provides insights into programmed genome rearrangement and vertebrate evolution. *Nat Genet* 50(2):270-277. <https://doi.org/10.1038/s41588-017-0036-1>.
- Specker JL, Kobuke L (1987) Seawater-acclimation and the thyroidal response to thyrotropin in juvenile coho salmon (*Oncorhynchus kisutch*). *J Exp Zool* 241: 327-332. <https://doi.org/10.1002/jez.1402410307>
- Specker JL, Schreck CB (1982) Changes in plasma corticosteroids during smoltification of coho salmon, *Oncorhynchus kisutch*. *Gen Comp Endocrinol* 46:53–58. [https://doi.org/10.1016/0016-6480\(82\)90162-9](https://doi.org/10.1016/0016-6480(82)90162-9)
- Sturm A, Bury N, Dengreville L, et al (2005) 11-Deoxycorticosterone is a potent agonist of the rainbow trout (*Oncorhynchus mykiss*) mineralocorticoid receptor. *Endocrinology* 146:47–55. <https://doi.org/10.1210/en.2004-0128>
- Sugimoto A, Oka K, Sato R, et al (2016) Corticosteroid and progesterone transactivation of mineralocorticoid receptors from Amur sturgeon and tropical gar. *Biochem J* 473:3655–3665. <https://doi.org/10.1042/BCJ20160579>
- Takabe S, Inokuchi M, Yamaguchi Y, Hyodo S (2016) Distribution and dynamics of branchial ionocytes in houndshark reared in full-strength and diluted seawater environments. *Comp Biochem Physiol A* 198:22-32. <https://doi.org/10.1016/j.cbpa.2016.03.019>
- Takahashi H, Sakamoto T (2013) The role of “mineralocorticoids” in teleost fish: relative importance of glucocorticoid signaling in the osmoregulation and “central” actions of mineralocorticoid receptor. *Gen Comp Endocrinol* 181:223-228. <https://doi.org/10.1016/j.ygcen.2012.11.016>
- Takei Y (2021) The digestive tract as an essential organ for water acquisition in marine teleosts: lessons from euryhaline eels. *Zoological Lett* 7(1):10. <https://doi.org/10.1186/s40851-021-00175-x>
- Takei Y, McCormick SD (2013) Hormonal control of fish euryhalinity. In: McCormick SD, Brauner CJ, Farrell AP (eds) *Fish Physiology Vol. 32 Euryhaline Fishes*. Academic Press, Amsterdam, pp 69-123
- Takei Y, Hiroi J, Takahashi H, Sakamoto T (2014) Diverse mechanisms for body fluid regulation in teleost fishes. *Am J Physiol Regul Integr Comp Physiol* 307(7):R778-792. <https://doi.org/10.1152/ajpregu.00104.2014>
- Takvam M, Denker E, Gharbi N, et al (2021) Sulfate homeostasis in Atlantic salmon is associated with differential regulation of salmonid-specific paralogs in gill and kidney. *Physiol Rep* 9(19):e15059. <https://doi.org/10.14814/phy2.15059>
- Tang CH, Lee TH (2011) Ion-deficient environment induces the expression of basolateral chloride channel, CIC-3-like protein, in gill mitochondrion-rich cells for chloride uptake of the tilapia *Oreochromis mossambicus*. *Physiol Biochem Zool* 84(1):54-67. <https://doi.org/10.1086/657161>
- Tingaud-Sequeira A, Calusinska M, Finn RN, et al (2010) The zebrafish genome encodes the largest vertebrate repertoire of functional aquaporins with dual paralogy and substrate specificities similar to mammals. *BMC Evol Biol* 10:38. <https://doi.org/10.1186/1471-2148-10-38>
- Tipsmark CK (2018) Identification of FXYP protein genes in a teleost: tissue-specific expression and response to salinity change. *Am J Physiol Regul Integr Comp Physiol* 294(4):R1367-1378. <https://doi.org/10.1152/ajpregu.00454.2007>
- Tipsmark CK, Jørgensen C, Engelund M, et al (2009) Effects of cortisol, growth hormone and prolactin on gill claudin expression in Atlantic salmon. *Gen Comp Endocrinol* 163:270–277. <https://doi.org/10.1016/j.ygcen.2009.04.020>
- Tipsmark CK, Madsen SS (2009) Distinct hormonal regulation of Na<sup>+</sup>,K<sup>+</sup>-atpase genes in the gill of Atlantic salmon (*Salmo salar* L.). *J Endocrinol* 203:301–310. <https://doi.org/10.1677/JOE-09-0281>
- Tipsmark CK, Madsen SS, Seidelin M, et al (2002) Dynamics of Na<sup>+</sup>,K<sup>+</sup>,2Cl<sup>-</sup> cotransporter and Na<sup>+</sup>,K<sup>+</sup>-ATPase expression in the branchial epithelium of brown trout (*Salmo trutta*) and Atlantic salmon (*Salmo salar*). *J Exp Zool* 293:106–118. <https://doi.org/10.1002/jez.10118>
- Tipsmark CK, Baltzegar DA, Ozden O, et al (2008a) Salinity regulates claudin mRNA and protein expression in the teleost gill. *Am J Physiol Regul Integr Comp Physiol* 294(3):R1004-1014. <https://doi.org/10.1152/ajpregu.00112.2007>
- Tipsmark CK, Kiilerich P, Nilsen TO, et al (2008b) Branchial expression patterns of claudin isoforms in Atlantic salmon during seawater acclimation and smoltification. *Am J Physiol Regul Integr Comp Physiol* 294(5):R1563-1574. <https://doi.org/10.1152/ajpregu.00915.2007>
- Tipsmark CK, Sørensen KJ, Madsen SS (2010) Aquaporin expression dynamics in osmoregulatory tissues of Atlantic salmon during smoltification and seawater acclimation. *J Exp Biol* 213(3):368-379. <https://doi.org/10.1242/jeb.034785>

- Tipsmark CK, Breves JP, Seale AP, et al (2011) Switching of Na<sup>+</sup>, K<sup>+</sup>-ATPase isoforms by salinity and prolactin in the gill of a cichlid fish. *J Endocrinol* 209(2):237-244. <https://doi.org/10.1530/JOE-10-0495>
- Tomy S, Chang YM, Chen YH, et al (2009) Salinity effects on the expression of osmoregulatory genes in the euryhaline black porgy *Acanthopagrus schlegelii*. *Gen Comp Endocrinol* 161(1):123-132. <https://doi.org/10.1016/j.ygcen.2008.12.003>
- Trayer V, Hwang PP, Prunet P, Thermes V (2013) Assessment of the role of cortisol and corticosteroid receptors in epidermal ionocyte development in the medaka (*Oryzias latipes*) embryos. *Gen Comp Endocrinol* 194:152-161. <https://doi.org/10.1016/j.ygcen.2013.09.011>
- Tse WK, Au DW, Wong CK (2006) Characterization of ion channel and transporter mRNA expressions in isolated gill chloride and pavement cells of seawater acclimating eels. *Biochem Biophys Res Commun* 346(4):1181-1190. <https://doi.org/10.1016/j.bbrc.2006.06.028>
- Tse WK, Au DW, Wong CK (2007) Effect of osmotic shrinkage and hormones on the expression of Na<sup>+</sup>/H<sup>+</sup> exchanger-1, Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> cotransporter and Na<sup>+</sup>/K<sup>+</sup>-ATPase in gill pavement cells of freshwater adapted Japanese eel, *Anguilla japonica*. *J Exp Biol* 210(Pt 12):2113-2120. <https://doi.org/10.1242/jeb.004101>
- Vialle RA, de Souza JES, Lopes KP, et al (2018) Whole genome sequencing of the pirarucu (*Arapaima gigas*) supports independent emergence of major teleost clades. *Genome Biol Evol* 10(9):2366-2379. <https://doi.org/10.1093/gbe/evy130>
- Wang PJ, Lin CH, Hwang HH, Lee TH (2008) Branchial FXD protein expression in response to salinity change and its interaction with Na<sup>+</sup>/K<sup>+</sup>-ATPase of the euryhaline teleost *Tetraodon nigroviridis*. *J Exp Biol* 211(Pt 23):3750-3758. <https://doi.org/10.1242/jeb.018440>
- Wang YF, Tseng YC, Yan JJ, et al (2009) Role of SLC12A10.2, a Na-Cl cotransporter-like protein, in a Cl uptake mechanism in zebrafish (*Danio rerio*). *Am J Physiol Regul Integr Comp Physiol* 296(5):R1650-1660. <https://doi.org/10.1152/ajpregu.00119.2009>
- Wang YF, Yan JJ, Tseng YC, et al (2015) Molecular physiology of an extra-renal Cl<sup>-</sup> uptake mechanism for body fluid Cl<sup>-</sup> homeostasis. *Int J Biol Sci* 11(10):1190-1203. <https://doi.org/10.7150/ijbs.11737>
- Watanabe S, Kaneko T, Aida K (2005) Aquaporin-3 expressed in the basolateral membrane of gill chloride cells in Mozambique tilapia *Oreochromis mossambicus* adapted to freshwater and seawater. *J Exp Biol* 208(Pt 14):2673-2682. <https://doi.org/10.1242/jeb.01684>
- Watanabe S, Itoh K, Kaneko T (2016) Prolactin and cortisol mediate the maintenance of hyperosmoregulatory ionocytes in gills of Mozambique tilapia: exploring with an improved gill incubation system. *Gen Comp Endocrinol* 232:151-159. <https://doi.org/10.1016/j.ygcen.2016.04.024>
- Weisbart M, Chakraborti PK, Gallivan G, Eales JG (1987) Dynamics of cortisol receptor activity in the gills of the brook trout, *Salvelinus fontinalis*, during seawater adaptation. *Gen Comp Endocrinol* 68:440-448.
- Weng CF, Lee TH, Hwang PP (1997) Immune localization of prolactin receptor in the mitochondria-rich cells of the euryhaline teleost (*Oreochromis mossambicus*) gill. *FEBS Lett* 405(1):91-94. [https://doi.org/10.1016/s0014-5793\(97\)00162-2](https://doi.org/10.1016/s0014-5793(97)00162-2)
- Whitehead A, Roach JL, Zhang S, Galvez F (2011) Genomic mechanisms of evolved physiological plasticity in killifish distributed along an environmental salinity gradient. *Proc Natl Acad Sci USA* 108(15):6193-6198. <https://doi.org/10.1073/pnas.1017542108>
- Wongdee K, Charoenphandhu N (2013) Regulation of epithelial calcium transport by prolactin: from fish to mammals. *Gen Comp Endocrinol* 181:235-240. <https://doi.org/10.1016/j.ygcen.2012.07.006>
- Wright PA, Wood CM (2015) Regulation of ions, acid-base, and nitrogenous wastes in elasmobranchs. In: Shadwick RE, Farrell AP, Brauner CJ (eds) *Fish Physiology*. Academic Press, pp 279-345.
- Wu CY, Lee TH, Tseng DY (2023) Mineralocorticoid receptor mediates cortisol regulation of ionocyte development in tilapia (*Oreochromis mossambicus*) *Fishes* 8(6):283. <https://doi.org/10.3390/fishes8060283>
- Xu B, Miao H, Zhang P, Li D (1997) Osmoregulatory actions of growth hormone in juvenile tilapia (*Oreochromis niloticus*). *Fish Physiol Biochem* 17:295-301. <https://doi.org/10.1023/A:1007750022878>
- Yada T, Miyamoto K, Miura G, Munakata A (2014) Seasonal changes in gene expression of corticoid receptors in anadromous and non-anadromous strains of rainbow trout *Oncorhynchus mykiss*. *J Fish Biol* 85:1263-1278. <https://doi.org/10.1111/jfb.12521>

- Yamaguchi Y, Ikeba K, Yoshida MA, Takagi W (2023) Molecular basis of the unique osmoregulatory strategy in the inshore hagfish, *Eptatretus burgeri*. *Am J Physiol Regul Integr Comp Physiol*. <https://doi.org/10.1152/ajpregu.00166.2023>.
- Yamaguchi, K., Hara, Y., Tatsumi, K., et al (2020) Inference of a genome-wide protein coding gene set of the inshore hagfish *Eptatretus burgeri*. *bioRxiv* 2020.07.24.218818. <https://doi.org/10.1101/2020.07.24.218818>
- Yan JJ, Hwang PP (2019) Novel discoveries in acid-base regulation and osmoregulation: a review of selected hormonal actions in zebrafish and medaka. *Gen Comp Endocrinol* 277:20-29. <https://doi.org/10.1016/j.ygcen.2019.03.007>
- Yao K, Niu PD, Gac FL, Bail PYL (1991) Presence of specific growth hormone binding sites in rainbow trout (*Oncorhynchus mykiss*) tissues: characterization of the hepatic receptor. *Gen Comp Endocrinol* 81:72–82. [https://doi.org/10.1016/0016-6480\(91\)90126-Q](https://doi.org/10.1016/0016-6480(91)90126-Q)
- Young G, Björnsson BT, Prunet P, et al (1989) Smoltification and seawater adaptation in coho salmon (*Oncorhynchus kisutch*): plasma prolactin, growth hormone, thyroid hormones, and cortisol. *Gen Comp Endocrinol* 74:335–345. [https://doi.org/10.1016/s0016-6480\(89\)80029-2](https://doi.org/10.1016/s0016-6480(89)80029-2)
- Yu D, Ren Y, Uesaka M, et al (2023) Hagfish genome illuminates vertebrate whole genome duplications and their evolutionary consequences. *bioRxiv* 2023.04.08.536076. <https://doi.org/10.1101/2023.04.08.536076>
- Zhou B, Kelly SP, Janowski JP, Wood CM (2003) Effects of cortisol and prolactin on Na<sup>+</sup> and Cl<sup>-</sup> transport in cultured branchial epithelia from FW rainbow trout. *Am J Physiol Regul Integr Comp Physiol* 285(6):R1305-1316. <https://doi.org/10.1152/ajpregu.00704.2002>

## Figure legends

**Figure 1.** Schematic diagrams of “Type II” and “Type III” ionocytes in Mozambique tilapia showing the stimulatory (arrows with a “+”) effects of prolactin (Prl) (see text for citations). Nka- $\alpha$ 1a and Clc2c are included in these models based upon the expression of their associated gene transcripts; however, they have yet to be definitively assigned to tilapia ionocytes. Apical and basolateral sides are presented at the top and bottom of cells, respectively. Abbreviations: Aqp3: aquaporin 3; Clc2c: Clc family Cl<sup>-</sup> channel 2c; Ncc2: Na<sup>+</sup>/Cl<sup>-</sup> cotransporter 2; Nka: Na<sup>+</sup>/K<sup>+</sup>-ATPase; Prl: prolactin.


**Figure 2.** Schematic diagrams of FW (freshwater)- and SW (seawater)-type ionocytes in mummichogs showing the stimulatory (arrows with a “+”) and inhibitory (blocked lines with a “-”) effects of prolactin (Prl) (see text for citations). Where Cl<sup>-</sup> transport is indicated with a question mark, a pathway is presumed to exist but remains uncharacterized. Apical and basolateral sides are presented at the top and bottom of cells, respectively. Abbreviations: Aqp3: aquaporin 3; Cftr1: cystic fibrosis transmembrane conductance regulator 1; Cldn10f: claudin 10f; Ncc2: Na<sup>+</sup>/Cl<sup>-</sup> cotransporter 2; Nka: Na<sup>+</sup>/K<sup>+</sup>-ATPase; Nkcc1: Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> cotransporter 1; Prl: prolactin; TJ: tight-junction. Figure adapted from Breves et al. (2022).

**Figure 3.** Schematic diagram of SW (seawater)-type ionocytes showing the stimulatory (arrows with a “+”) and inhibitory (blocked lines with a “-”) effects of cortisol (Cort) (see text for citations). Apical and basolateral sides are presented at the top and bottom of


1821 cells, respectively. Abbreviations: Aqp3: aquaporin 3; Cftr1: cystic fibrosis  
1822 transmembrane conductance regulator 1; Cldn10s: claudin 10 isoforms; Cort: cortisol;  
1823 Nka:  $\text{Na}^+/\text{K}^+$ -ATPase; Nkcc1:  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  cotransporter 1; TJ: tight-junction.



**Fig. 1**



**Fig. 2**



**Fig. 3**

