

PROTOCOL

Aim-Tconv Human HMP

Version 1.1.7


FOR RESEARCH USE ONLY
STORE IN 2-8°C

Human T Cell Activator

CATALOG ITEMS: ATC01-0050H

Description

Aim-Tconv hydrogel microparticles (HMP) are designed for robust activation of human T cells using a feeder-free culturing approach. For any downstream processing (e.g. counting, flow cytometry) or earlier removal of HMP, please remove using the HMP digesting buffer provided in the kit.



Components

A suspension of hydrogel microparticles (HMP) made of chemically crosslinked dextran. HMPs sized 15 μm were coated with phospholipid bilayer, with membrane docked human T cell activating signal panels. Each kit contains:

- ◆ 1 vial of Aim-Tconv in PBS at 4×10^7 beads / mL, with 0.5% P/S
ATC01-0010H: 0.1 mL, total HMP number = 4×10^6
ATC01-0050H: 0.5 mL, total HMP number = 2×10^7
- ◆ 1 vial of HMP digesting buffer (100X, 0.5 mL), contains dextranase in PBS.

Stability and Storage

- ◆ Shipped with ice, keep product refrigerated (2-8°C)
- ◆ Stable at 4°C for 12 months
- ◆ Contents are sterile in unopened tube
- ◆ Do not subject product to freezing, high temperatures (>40°C)

Other Required Materials

- Cryopreserved or freshly isolated human PBMC, or isolated CD3⁺ T Cells
- RPMI 1640 supplemented with 2 mM L-glutamine (or equivalent)
- Heat inactivated Fetal Bovine Serum (FBS)
- Penicillin/streptomycin
- Recombinant human IL-2 (rIL-2)
- 2-Mercaptoethanol
- Cell culture vessels
- Humidified CO₂ incubator or bioreactor

Activation Protocol

Medium preparation

RPMI, 10% FBS, 1% Pen/Strep, 20~50 μM 2-Mercaptoethanol, 30~100 U/mL rIL-2
Or any other compatible culture media.

HMP preparation

Resuspend HMP in the vial by vortexing for 30 seconds. Open HMP vial and HMP lysis buffer only in sterile environment to avoid contamination. Calculate the desired HMP seeding density per well and aliquot the HMP into each well accordingly. Rather than recommending a fixed bead-to-cell ratio, we provide a validated seeding density range for reference. We also encourage experienced users to optimize seeding conditions based on their specific experimental requirements. The HMP seeding principle is to balance the interaction probability between HMP and target cells while leaving sufficient space for cells to grow.

Recommendations (seeding density range validated)

- ◆ Isolated T cell, or PBMC seeding density : $1.5 \times 10^4 \sim 2.5 \times 10^5$ cell per cm^2
- ◆ HMP seeding density : $8 \times 10^4 \sim 1.6 \times 10^5$ HMP per cm^2

Plate	Area (cm^2)	Cell/well	Cell/ cm^2	HMP/well	HMP/ cm^2	HMP Vol.
96 well	0.32	$5.0 \times 10^3 \sim 8.0 \times 10^4$	$1.5 \times 10^4 \sim 2.5 \times 10^5$	$2.5 \times 10^4 \sim 5.0 \times 10^4$	$8.0 \times 10^4 \sim 1.6 \times 10^5$	0.6~1.3 μL
48 well	0.95	$2.4 \times 10^4 \sim 2.4 \times 10^5$		$7.6 \times 10^4 \sim 1.5 \times 10^5$		1.9~3.8 μL
24 well	1.9	$4.8 \times 10^4 \sim 4.8 \times 10^5$		$1.5 \times 10^5 \sim 3.0 \times 10^5$		3.8~7.6 μL
6 well	9.5	$2.4 \times 10^5 \sim 2.4 \times 10^6$		$7.6 \times 10^5 \sim 1.5 \times 10^6$		19~38 μL
T25 flask	25	$6.3 \times 10^5 \sim 6.3 \times 10^6$		$2.0 \times 10^6 \sim 4.0 \times 10^6$		50~100 μL
T75 flask	75	$1.9 \times 10^6 \sim 1.9 \times 10^7$		$6.0 \times 10^6 \sim 1.2 \times 10^7$		150~300 μL

Cell seeding

Aliquot resuspended PBMC or isolated T cells. Gently mix cells and HMP by pipetting up/down 3 to 5 times ensure HMP and cells are evenly distributed under microscope.

T cell activation

Incubate in a humidified 5% CO₂ incubator at 37°C. Monitor T cell morphology and confluency, perform half media replenishment every other day. **DO NOT disturb** HMP-cell interaction in the first 4 days.

Restimulation

T cell growth typically slows after day 10 post-initial activation. Users may consider restimulation to promote continued cell expansion. Restimulating with Aim-Tconv HMP minimizes activation-induced exhaustion through its mild yet effective stimulation signal. We recommend using 8×10⁴ HMP per cm² for restimulation.

HMP cleanup

HMP will self-degrade through hydrolysis after approximately 8–10 days in serum consisting RPMI1640 media. Since the hydrolysis rate is pH-dependent, the actual degradation time may vary depending on the culture medium used.

Alternatively, HMP can be rapidly degraded enzymatically by adding HMP digesting buffer (included in the kit) directly to the culture medium at 1× final concentration. After incubating at 37°C for 30 minutes, the HMP will be fully degraded.

Important Note:

- ❖ HMP digesting buffer contains dextranase, which degrades the HMP core. Do not use this buffer if you plan to restimulate cells, as residual enzyme will degrade newly added HMP unless thoroughly washed out.
- ❖ It is recommended to degrade HMP before cryopreservation. Wash the cell pellet at least twice before cryopreservation to remove residual dextranase.

Cell Phenotype Characterization

- ❖ Resting T cells: smaller in size, round shaped
- ❖ Activated T cells: larger in size, irregular

Things to Note

When seeding cells with HMPs

Ensure cells and HMP are evenly distributed to maximize interaction.

When you see cell cluster

Aim-Tconv HMP and T cells tend to aggregate in the well centre over time. Gently shake the culture plate to redistribute HMP and T cells. **Avoid unnecessary pipette mixing in the first 4 days**, disturbing the HMP-cell clustering will cause suboptimal cell growth.

When to add new medium or split?

Monitor cell growth periodically by performing cell sampling and counting periodically. Supplement fresh medium or pass the cells to new culture vessels when:

- ❖ Colour turns orange yellow (acidic, ~ pH 6.6)
- ❖ Cell grows to over 3 x 10⁶ cells /mL

When to restimulate

Day 8-10 after previous stimulation.



Aim-Tconv is part of our AimGel artificial cell line-up.



www.allegrowbiotech.com

Disclaimer

This product is intended for research use only and is not approved for human or animal diagnostic or therapeutic applications. Allegrow Biotech warrants the product to conform substantially to its published specifications at the time of delivery, under normal use and conditions, until the stated expiration date or, if none is provided, for one (1) year from delivery. The purchaser is solely responsible for determining the suitability of the product for specific applications and for obtaining any necessary licenses. Allegrow Biotech disclaims all other warranties, express or implied, including merchantability or fitness for a particular purpose, and shall not be liable for incidental or consequential damages arising from product use.

Copyright © 2025 Allegrow Biotech. All rights reserved. Trademarks are the property of Allegrow Biotech or their respective owners.