

CORIANDER MEDICAL CLINIC, WINCHESTER

Birth (high-risk): Hepatitis B

2 months

- DTaP-IPV-Hib (Pentacel®)
- Pneumococcal (Pevnar 13®)
- Rotavirus (Rotarix®)

4 months

- DTaP-IPV-Hib (Pentacel®)
- Pneumococcal (Pevnar 13®)
- Rotavirus (Rotarix®)

6 months

- DTaP-IPV-Hib (Pentacel®)

12 months

- MMR (MMR II®)
- Meningococcal C (NeisVac-C®)
- Pneumococcal (Pevnar 13®)

15 months

- Varicella (Varivax®)

18 months

- DTaP-IPV (Adacel-Polio®)

4-6 years

- DTaP-IPV (Quadrace®)

Grade 7 (~12 yrs)

- HPV (2-dose, Gardasil® 9)
- Hepatitis B (2-dose)

14-16 yrs

- Meningococcal ACYW-135 (Nimenrix®)

ADULTS

- Td booster every 10 years
- One adult booster should be **Tdap**

Pregnancy (each pregnancy)

- Tdap at **27-32 weeks**

Age ≥65 years

- Pneumovax® 23
- Influenza (every year)

Risk-based (any age)

- Influenza • COVID-19 • Hepatitis A • Meningococcal B

Shingles

- Shingrix® (publicly funded ≥65 yrs)

RSV

Abrysvo: late pregnancy (32-36 wks) for passive immunity, and
Beyfortus/Nirsevimab: birth to 8mo