



Date: _____ Patient Name: _____
 DOB: _____ Address: _____
 City: _____ State: _____ Phone: _____ Allergies: _____
 Call When Ready Text Message When Ready Delivery Mail Out

- Testosterone in Vanishing Light Topical Cream**
(circle strength) 1% or 2%
 Qty: 9 ml
 Sig: Apply 0.1-0.3 ml once daily in the AM. Start at 0.1 ml daily for 2 weeks, then 0.2 ml for 2 weeks, then 0.3 ml daily if tolerated well.

- Libido #1 Theophylline 3%/Arginine(L) 6%/Sildenafil 1% Cream**
 Qty: # 10 ml
 Sig: Apply a pea-sized amount topically to clitoral area 5-30 min prior to intercourse as directed.

- Libido #2 Theophylline 3%/Arginine(L) 6%/Menthol 0.025%/Nifedipine 0.2% Cream**
 Qty: # 10 ml
 Sig: Apply a pea-sized amount topically to clitoral area 5-30 min prior to intercourse as directed.

- Libido #3 [Raspberry Flavored] Arginine 2%/Naltrexone 0.7% Gel**
 Qty: # 10 gm
 Sig: Apply a pea-sized amount topically to clitoral area 5-30 min prior to intercourse as directed.

- Oxytocin 120 U/gm Vaginal Gel**
(Mucolox™/Versabase®)
 Qty: # 45 gm
 Sig: Apply a pea-sized amount topically to clitoral area 5-30 min prior to intercourse as directed.

- Oxytocin Nasal Spray 6 units per actuation**
 Qty: # 15 ml - Each spray delivers 0.1 ml (6 Units)
 Sig: Instill 2 sprays in each nostril as needed

Refills: 1 2 3 4 5 PRN

 Healthcare Provider Signature:

Print Name: _____ Agent sending: _____
 NPI: _____ DEA: _____

Clinic Name: _____
 Clinic Address: _____
 Clinic Phone/Fax: _____

