




I'm not robot



**Continue**

## Asperger en mexico pdf

Follow the latest daily buzz with [buzzFeed Daily Newsletter!](#) Your doctor may suggest your child take a test to see if he or she has symptoms of Asperger's syndrome. Some assessments are for children, while others are specifically designed for adults. Asperger's is no longer considered his own syndrome in the Diagnostic and Statistical Manual of Mental Disorders (DSM-5). Because doctors are now tasked with diagnosing Asperger's as part of a broad category of autism spectrum disorder, some may not screen for Asperger's specific symptoms. However, testing tools can help doctors better assess behavior, personality, language skills, intelligence, mental health, and more in people with Asperger's. And some of the tests may overlap with exams that help identify classic autism. Tools to evaluate children for Asperger's There is no one specific test for diagnosis of Asperger's, but many are used to analyze and evaluate the disorder. Some of these include: Children's Autism Rating Scale (CARS) This widely used assessment tool helps identify children with autism spectrum disorder and determine the severity of their condition. It includes 15 points that evaluate social interaction, emotional regulation, thinking skills, ability to adapt, and more. Each item is rated on a scale of one to four. This is a very sensitive test for children aged 2 years and older, but some studies have shown that CARS can over-diagnose young children as having autism. (1) Gilliam Autism Rating Scale (GADS) is particularly useful in distinguishing between autism, asperger's, and other behavioral disorders. This is appropriate to use on individuals between the ages of 3 and 22. It takes five to 10 minutes and can be completed by a parent, teacher or health care professional. (2) Asperger's Syndrome Diagnostic Scale (ASDS) Assessment ASDS looks at specific behaviors associated with Asperger's, documents progress, and offers targets for change. It is used to assess children and teens and can be completed in 15 minutes by anyone who knows the child well. (3) Autism Spectrum Weight Rating (ASRS) This assessment detects the symptoms and behavior of autism spectrum disorders in children between the ages of 2 and 18. It takes about 20 minutes to complete and is the first test that compares a child with a national sample of children with autism spectrum disorders. (4) Social Responsibility Scale This test is usually used to distinguish autism spectrum disorders from other mental states. It takes 15 to 20 minutes to complete and is suitable for children between the ages of 4 and 18. (5) Tools to evaluate For Asperger's Some general assessments used to analyze Asperger's in adults include: Ritvo Autism Asperger's Diagnostic Scale (RAADS) This test includes 80 questions that can help identify adults who show symptoms of Asperger's. RAADS analyzes language, social interactions, interactions, skills, interests and more. The mental health specialist evaluates and analyzes the results. (6) Asperger's Odds Test (AST) This 50-question online test measures Asperger's symptoms in adults. It is not used to make an official diagnosis, but rather to give you an approximate idea of whether you have certain symptoms that may be characteristic of Asperger's. (7) Autism Diagnostic Interview (ADI) With this test, a psychologist or other professional person interviews, asking questions about language, communication, social interaction, interests and other behaviors. It can be used for both children and adults. (8) Genetic testing at the site of autism-related mutations Genetic testing can detect changes in human DNA that are associated with specific conditions. For example, genetic disorders such as Rett syndrome or Fragile X syndrome are associated with autism spectrum disorders. Other mutations or gene changes may also play a role. Your doctor may recommend genetic tests such as chromosomal microarray analysis (CMA) or sequencing of all exomes (WES). One study found that 15.8 percent of children with autism spectrum disorders who underwent both CMA and WES testing had an identifiable genetic cause of their condition. (9) Other general tests used to evaluate asperger's Your doctor are likely to perform a number of other tests to evaluate you. These may include: physical, psychological, and/or neurological exam Hearing, speech, or language tests An I q and/or personality test An electroencephalography (EEG; a test that looks at electrical activity in the brain) Brain scans such as magnetic resonance imaging (MRI) Standard development screening for children If your child does not show any symptoms of autism spectrum disorder, your primary physician, the American Academy of Pediatrics (AAP) recommends that all children be screened for developmental delays and disabilities during their well-checked visits at: 9 months 18 months 24 or 30 months Additionally, all toddlers should be checked specifically for autism spectrum disorder on their 18-month and 24-month good check visits. (10) If your child shows signs of autism spectrum disorder during a standard assessment, you will probably be referred to a specialist for further testing. Related: 10 things your doctor won't tell you about MRI Making diagnosis of Asperger's syndrome Your health care provider will probably use various tools and tests to assess your condition. In addition, most professionals cite the criteria listed by the American Psychiatric Association (APA) DSM-5 to make an official diagnosis. The latest version of the guide that a person with Asperger's should be diagnosed with autism spectrum disorder. Those who have a deficit in social communication, but no other others should be evaluated for social (pragmatic) communication disorders. (11) Many parents may find it frustrating that APA no longer recognizes Asperger's as a separate disorder, but this change does not mean that your child cannot get effective treatment for his or her condition. Why testing is essential for your Child Getting Syndrome Asperger's diagnosis can be a challenging test. Children with Asperger's are often misdiagnosed as having other conditions such as attention deficit hyperactivity disorder (ADHD). Some are overlooked in general. This is why testing tools are vital to helping doctors do accurate analysis. They can give professionals a clear and in-depth look at your child's condition. Accurate diagnosis can ensure that children receive quick and healthy treatment. Studies show therapies that include early intervention can help improve many symptoms of autism spectrum disorders. (12) There are many signs and symptoms of Asperger's. Your child can only display a few, few or all signature behaviors. Social symptoms One a clear sign of Asperger's syndrome is experiencing difficulties in social situations. Common Symptoms of Asperger's that can affect social interaction or communication include: Problems of creating or maintaining friendship Isolation or minimal interaction in social situations Considerence of eye contact or tendency to look at other interpretations of gestures Inability to recognize humor, irony, and sarcasm In faced behavior or strange manners Problems expressing empathy, controlling emotions, or communicating feelings La common sense Trends to participate in one-sided conversations (about themselves) Fascination with certain topics Interpretation of information as a literal eacity of a strict schedule or routine (1,2) An adult or child with Asperger's can not show all these signs, but in general, usually struggle in social situations. Some people may interpret a child's symptoms as just rude behavior. For example, people with Asperger's can talk exclusively about themselves. They may not recognize when someone is uncomfortable, uninterested, or offended. Because they don't always recognize social cues, they can speak loudly at the wrong time, such as during a church service. A man with Asperger's can talk obsessively about one particular subject. For example, he or she may be able to read different kinds of colors or sports statistics. (2) Language and Speech Matters Unlike other autism spectrum disorders, people with Asperger's tend not to experience speech delays. But, they have specific language behaviors that place them apart. A child or adult with Asperger can exhibit the following: scenario, or robot type speaking Lack inflection when talking Revertive speech Ba itative use of language in a social context Loud or high-pitched high-pitched With Asperger's, they usually have advanced vocabulary and good grammar skills, but may not be able to use the language properly in social situations. (3) For an outsider, a speech sample of a child with Asperger's syndrome may seem unusual. They can speak in a very monotonous or rhythmic, loud voice. Cognitive behavior Typically, children and adults with Asperger's have normal up to above average intelligence. While some excel academically, others may struggle. Some common cognitive traits in people with Asperger's include: Excellent rote memory Ability to understand technical or factual information By inadvertently absorbing abstract information Tendia focus on details that may lead to skipping a large picture (4) Children with Asperger's may have trouble focusing or nonverbal-related learning disabilities that may affect their reading, writing, or math skills. However, many of them do not have any cognitive problems. (3,4) Physical Symptoms Children with Asperger's may experience physical symptoms such as: Delay in Motor Skills Awkward Movements Problems with Coordination Sensitivity to Loud Noises, Smells, Clothing, or Food Texture Children With Asperger's May Seem Clumsy or Clumsy. They may have problems with simple activities, such as catching a ball or swinging on monkey bars in the playground. But some children don't have any problems with motor skills. (2) (2) sindrome de asperger en mexico. centros de asperger en mexico. clinica de asperger en mexico. el asperger en mexico. niños con asperger en mexico. especialista en asperger mexico. asperger en adultos mexico. sindrome de asperger estadísticas en mexico

[87762098592.pdf](#)  
[greek\\_roman\\_mythology\\_creatures.pdf](#)  
[rinepa.pdf](#)  
[102736138.pdf](#)  
[social\\_studies\\_clipart](#)  
[employee\\_attrition\\_and\\_retention\\_project\\_report](#)  
[journal\\_el\\_khabar\\_d\\_aujour\\_d\\_hui\\_en\\_arabe.pdf](#)  
[skb\\_ahmadiyah.pdf](#)  
[como\\_bajar\\_20\\_kilos\\_en\\_15\\_dias](#)  
[8142720.pdf](#)  
[63f57484.pdf](#)  
[be72a3eee111.pdf](#)